

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

5-2003

Alpha-Toxin Is Required for Biofilm Formation by *Staphylococcus aureus*

Nicky C. Caiazza
Dartmouth College

George A. O'Toole
Dartmouth College

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Bacteriology Commons](#), and the [Medical Microbiology Commons](#)

Dartmouth Digital Commons Citation

Caiazza, Nicky C. and O'Toole, George A., "Alpha-Toxin Is Required for Biofilm Formation by *Staphylococcus aureus*" (2003). *Dartmouth Scholarship*. 1109.
<https://digitalcommons.dartmouth.edu/facoa/1109>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Alpha-Toxin Is Required for Biofilm Formation by *Staphylococcus aureus*

Nicky C. Caiazza and G. A. O'Toole*

Department of Microbiology and Immunology, Dartmouth Medical School, Hanover, New Hampshire

Received 9 May 2002/Accepted 24 February 2003

***Staphylococcus aureus* is a common pathogen associated with nosocomial infections. It can persist in clinical settings and gain increased resistance to antimicrobial agents through biofilm formation. We have found that alpha-toxin, a secreted, multimeric, hemolytic toxin encoded by the *hla* gene, plays an integral role in biofilm formation. The *hla* mutant was unable to fully colonize plastic surfaces under both static and flow conditions. Based on microscopy studies, we propose that alpha-hemolysin is required for cell-to-cell interactions during biofilm formation.**

Biofilms are surface-associated, sessile bacterial communities. A mature biofilm is formed when planktonic cells initially colonize a surface, aggregate and/or grow into multicellular colonies, and embed themselves in an exopolysaccharide matrix. *Staphylococcus aureus* is capable of biofilm formation, which increases its persistence and boosts its levels of antimicrobial resistance (5), and biofilms of this organism have been observed on surfaces ranging from intravascular catheters to pacemaker leads (17, 18). Genetic analyses of staphylococci have shown that the progression of biofilm development consists of two steps: initial cell-to-surface interactions followed by cell-to-cell interactions (9). Recent reports have shown autolysin (10), teichoic acids (8), and surface proteins such as Bap to be integral to the initial stages of colonization (4). The *ica* locus, which is required for the synthesis of the polysaccharide intracellular adhesin (PIA), plays a role in subsequent cell-to-cell interactions (3, 14).

The accessory gene regulator (*agr*) is a two-component regulatory system in *S. aureus* that has been implicated in biofilm formation—an *agr* mutant is a hyper-biofilm-forming strain (22). To study biofilms of *S. aureus*, we took the approach of examining known downstream targets regulated by the *agr* system and determining their impact on biofilm formation. We show that one of these targets, alpha-hemolysin, a 34-kDa protein that causes host cell lysis by heptamerizing upon insertion into eukaryotic cell membranes, plays a role in biofilm formation (21, 23). Mutants defective in alpha-hemolysin production failed to form biofilms under both static and flow conditions, and strains lacking alpha-hemolysin have an apparent defect in cell-to-cell interactions.

Figure 1A shows the results of a biofilm assay wherein bacteria were grown at 37°C in tryptic soy broth (TSB) and 0.2% glucose for 8 h, as described by Heilmann et al. (9, 11). The level of bacterial adhesion, as quantified by crystal violet staining, is ~3-fold lower for the *hla* mutant than for the wild type (strains are described in Table 1). The alpha-hemolysin-defi-

cient strain was also defective for biofilm formation when compared to the wild type at 16 h (data not shown). Plasmid pDU1212 contains a wild-type copy of the *hla* gene, and when this plasmid is introduced into the *hla::erm* strain, biofilm formation is induced to a level above that of even the wild-type strain (Fig. 1B), whereas the vector control pNC1 has no effect




FIG. 1. The *hla* mutant is defective in biofilm formation. (A) Biofilm formation phenotypes. Biofilm formation by the wild type (WT; *S. aureus* 8325-4) and a *hla::erm* mutant (*S. aureus* DU1090) was quantitated. Crystal violet was used to stain cells adhering to polystyrene after 8 h of growth at 37°C. (B) Complementation of the *hla::erm* mutant. Biofilms formed on polystyrene (8 h at 37°C) were analyzed for an *hla::erm* mutant harboring a vector control plasmid (pNC1) or a plasmid providing a wild-type copy of the *hla* gene (pDU1212 [*hla*⁺]).

* Corresponding author. Mailing address: Department of Microbiology & Immunology, Rm. 202 Vail Building, Dartmouth Medical School, Hanover, NH 03755. Phone: (603) 650-1248. Fax: (603) 650-1318. E-mail: georgeo@Dartmouth.edu.

TABLE 1. Strains and plasmids

Strain or plasmid	Description	Reference
<i>S. aureus</i> strains		
8325-4		19
DU1090	Strain 8325-4 with <i>hla::erm</i>	20
DU1090/pDU1212	Strain 8325-4 with <i>hla::erm</i> , harboring pDU1212 (Ap ^r Cm ^r / <i>hla</i> ⁺)	6
DU1090/pNC1	Strain 8325-4 with <i>hla::erm</i> , harboring pNC1 (Ap ^r Cm ^r)	This study
113		3
113 <i>ica::Tc</i>	Defective in production of PIA	3
113 <i>ica::Tc</i> /pNC1	Cm ^r	This study
113 <i>ica::Tc</i> /pDU1212	Cm ^r	This study
Plasmids		
pDU1212	pBR322 carrying <i>hla</i> ⁺ and Cm ^r	6
pNC1	Empty vector control; pBR322 carrying 3.0-kb <i>Hind</i> III fragment from pDU1212 encoding Cm ^r	This study

on biofilm formation. It has been shown previously that the supernatant of *S. aureus* DU1090/pDU1212 (*hla*⁺) contains 2.5- and 110-fold more hemolytic activity (in hemolytic units per milliliter) than wild-type and *hla::erm* strains, respectively (1). Thus, the level of alpha-hemolysin may correlate with the level of biofilm formation.

PIA, encoded by the *ica* genes, has been shown to be required for biofilm formation by *S. aureus* (3, 14, 15); therefore, we investigated PIA production in the wild-type and *hla::erm* strains. PIA was extracted from cells grown in TSB supplemented with 0.2% glucose (the medium used for biofilm as-

says), serial twofold dilutions were spotted onto nitrocellulose, and Western blotting was performed as previously described by Cramton et al. (3) by using antibody to PIA/PNAG [β (1-6)-*N*-acetylglucosamine] (16). A wild-type PIA-producing strain (113) and an isogenic *ica* mutant (113 *ica::Tc*) served as controls. No difference in the levels of PIA production between the wild-type and alpha-toxin mutant strains was observed (data not shown). Furthermore, in 10 clinical *S. aureus* strains analyzed (22), no correlation between PIA production and alpha-hemolysis was observed. We also investigated the ability of a multicopy dose of *hla* (plasmid pDU1212) to rescue the




FIG. 2. Direct visualization of attachment phenotypes. Bacteria were inoculated onto 24-well plates, incubated for 8 h at 37°C, and then analyzed by phase-contrast microscopy. Dark areas are the adherent bacteria, and the light grayish regions represent the surface of the 24-well plate. The magnification is $\times 1,050$. WT, wild type.




FIG. 3. Phenotypes of cells under flow conditions. Biofilms of the wild-type (WT; *S. aureus* 8325-4) and the *hla::erm* mutant (*S. aureus* DU1090) strains were grown in flow cell chambers. At the times indicated, biofilms were observed from a top-down perspective by using phase-contrast microscopy. In images from 4 and 8 h, light regions represent bacterial macrocolonies (indicated by the white arrows) and dark areas are the surface of the flow cell chamber. The magnification is $\times 675$ for images from 4 and 8 h. At 24 h, the images were recorded at an original magnification of $\times 230$. (They are shown at a magnification of $\times 173$.) In these images from 24 h, the very dense macrocolonies formed by the wild type appear as dark regions and the light areas define macrocolony borders or channels between the macrocolonies. Small clusters of cells, but no macrocolonies, were observed for the *hla* mutant at all time points.

biofilm formation defect of an *ica* mutant. Neither pDU1212 (*hla*⁺) nor the vector control (pNC1) had any effect on the biofilm formation phenotype of the *ica* mutant (data not shown).

To better understand the nature of the biofilm-deficient phenotype of the *hla::erm* mutant, phase-contrast microscopy was employed to observe and compare levels of surface attachment at 8 h in 24-well polystyrene plates (Costar, Corning, N.Y.) (Fig. 2). This assay was similar to the 96-well plate assay (described in references 9 and 11) with the exception that nonadherent cells were removed by aspiration. For the wild-type strain, microcolonies (dark regions) were found scattered evenly throughout the field of view but were not present in the fields of view for the *hla::erm* mutant and the vector control. The strain carrying the plasmid pDU1212 (*hla*⁺) in the *hla::erm* background exhibited more robust biofilm formation

than even the wild type—the entire surface was covered in a dense mass of microcolonies. Therefore, the crystal violet staining data presented in Fig. 1B correlates with the microscopy data presented in Fig. 2.

In a physiological setting, such as the surface of a catheter, biofilms may exist and persist under conditions of flow. To mimic these conditions in vitro, *S. aureus* biofilms were grown under conditions of constant flow (40 ml/h) by using 0.1 \times TSB as the growth medium in the flow cell system described by Christensen et al. (2). Overnight cultures of *S. aureus* were diluted 1:1,000 in 0.1 \times TSB, 300 μ l of diluted cells was injected into the flow cell chamber, and the cells were allowed to acclimate for 15 min before being subjected to flow. Figure 3A shows that by 4 h the wild type had attached to the surface of the flow chamber and begun to form large macrocolonies. After 8 h of constant flow, the wild-type macrocolonies became

larger and more numerous. In addition, the surface area between macrocolonies was completely covered by a monolayer of cells. By 24 h, wild-type macrocolonies had increased in size and density to the point of completely filling the flow chamber. The architecture of the wild-type biofilm at 24 h consisted of densely packed circular macrocolonies outlined by narrow, light regions that were the channels between the macrocolonies. In contrast to the wild type, the *hla::erm* mutant attached to the surface as a sparse monolayer, failed to exhibit macrocolony formation even at 24 h, and lacked any discernible architecture.

In this study, we show a role for alpha-hemolysin in *S. aureus* biofilm formation, and in particular, this toxin appears to be required for cell-to-cell interactions. We were initially surprised to find that a secreted toxin had such a dramatic impact on biofilm formation; however, other examples exist in which secreted toxins may play a role in biofilm formation (12, 13, 22). The fact that cells carrying a mutant allele of *hla* are capable of initially colonizing a surface but never organize into multicellular macrocolonies indicates a defect in cell-to-cell interactions. Based on the data presented in this study, we propose that alpha-hemolysin plays a role primarily in cell-to-cell interactions during biofilm formation.

Alpha-hemolysin is, in part, controlled by the *agr* system. It has been shown that an *agr* mutant produces less alpha-hemolysin but is a hyper-biofilm-forming strain (22). However, the *agr* system regulates a wide array of virulence factors, including those involved in surface binding and surface-associated virulence. Thus, even though alpha-hemolysin production is reduced in an *agr* mutant, other surface-associated virulence factors may be overexpressed, functionally compensating for the lack of alpha-hemolysin. Furthermore, in vivo studies of device-related infections have shown that alpha-hemolysin is not regulated by *agr* but that its expression is predominately controlled by the two-component regulator *sae* (7). Therefore, alpha-hemolysin may be produced in an *agr*-independent fashion when *S. aureus* colonizes in-dwelling devices in the biofilm mode of growth.

We thank A. Chueng and M. Palma for helpful advice and Richard J. O'Callaghan for sending strains DU1090 and DU1090/pDU1212. We also thank Jerry Pier for providing the PIA/PNAG antibodies and Cuong Vuong and Michael Otto for providing clinical *S. aureus* isolates.

This work was supported by grants from Microbia, Inc., by the American Cancer Society institutional research grant #IRG-82-003-17, and by the Pew Charitable Trusts to G.A.O. G.A.O. is a Pew Scholar in the Biomedical Sciences.

REFERENCES

- Bayer, A. S., M. D. Ramos, B. E. Menzies, M. R. Yeaman, A. J. Shen, and A. L. Cheung. 1997. Hyperproduction of alpha-toxin by *Staphylococcus aureus* results in paradoxically reduced virulence in experimental endocarditis: a host defense role for platelet microbicidal proteins. *Infect. Immun.* **65**: 4652–4660.
- Christensen, B. B., C. Sternberg, J. B. Andersen, R. J. Palmer, Jr., A. T. Nielsen, M. Givskov, and S. Molin. 1999. Molecular tools for study of biofilm physiology. *Methods Enzymol.* **310**:20–42.
- Cramton, S. E., C. Gerke, N. F. Schnell, W. W. Nicols, and F. Gotz. 1999. The intercellular adhesion (*ica*) locus is present in *Staphylococcus aureus* and is required for biofilm formation. *Infect. Immun.* **67**:5427–5433.
- Cucarella, C., C. Solano, J. Valle, B. Amorena, I. Lasa, and J. R. Penades. 2001. Bap, a *Staphylococcus aureus* surface protein involved in biofilm formation. *J. Bacteriol.* **183**:2888–2896.
- Cunningham, R., and J. Cheesbrough. 1992. Comparative activity of glycopeptide antibiotics against coagulase-negative staphylococci embedded in fibrin clots. *J. Antimicrob. Chemother.* **30**:321–326.
- Fairweather, N., S. Kennedy, T. J. Foster, M. Kehoe, and G. Dougan. 1983. Expression of a cloned *Staphylococcus aureus* alpha-hemolysin determinant in *Bacillus subtilis* and *Staphylococcus aureus*. *Infect. Immun.* **41**:1112–1117.
- Goerke, C., U. Fluckiger, A. Steinhuber, W. Zimmerli, and C. Wolz. 2001. Impact of the regulatory loci *agr*, *sarA* and *sae* of *Staphylococcus aureus* on the induction of alpha-toxin during device-related infection resolved by direct quantitative transcript analysis. *Mol. Microbiol.* **40**:1439–1447.
- Gross, M., S. E. Cramton, F. Gotz, and A. Peschel. 2001. Key role of teichoic acid net charge in *Staphylococcus aureus* colonization of artificial surfaces. *Infect. Immun.* **69**:3423–3426.
- Heilmann, C., C. Gerke, F. Perdreau-Remington, and F. Gotz. 1996. Characterization of Tn917 insertion mutants of *Staphylococcus epidermidis* affected in biofilm formation. *Infect. Immun.* **64**:277–282.
- Heilmann, C., M. Hussain, G. Peters, and F. Gotz. 1997. Evidence for autolysin-mediated primary attachment of *Staphylococcus epidermidis* to a polystyrene surface. *Mol. Microbiol.* **24**:1013–1024.
- Heilmann, C., O. Schweitzer, C. Gerke, N. Vanittanakom, D. Mack, and F. Gotz. 1996. Molecular basis of intercellular adhesion in the biofilm-forming *Staphylococcus epidermidis*. *Mol. Microbiol.* **20**:1083–1091.
- Kachlany, S. C., D. H. Fine, and D. H. Figurski. 2000. Secretion of RTX leukotoxin by *Actinobacillus actinomycetemcomitans*. *Infect. Immun.* **68**:6094–6100.
- Kachlany, S. C., P. J. Planet, M. K. Bhattacharjee, E. Kollia, R. DeSalle, D. H. Fine, and D. H. Figurski. 2000. Nonspecific adherence by *Actinobacillus actinomycetemcomitans* requires genes widespread in bacteria and archaea. *J. Bacteriol.* **182**:6169–6176.
- Mack, D., W. Fischer, A. Krokotsch, K. Leopold, R. Hartmann, H. Egge, and R. Laufs. 1996. The intracellular adhesion involved in biofilm accumulation of *Staphylococcus epidermidis* is a linear β -1,6-linked glucosaminoglycan: purification and structural analysis. *J. Bacteriol.* **178**:175–183.
- Mack, D., M. Nedelmann, A. Krokotsch, A. Schwarzkopf, J. Heesemann, and R. Laufs. 1994. Characterization of transposon mutants of biofilm-producing *Staphylococcus epidermidis* impaired in the accumulative phase of biofilm production: genetic identification of a hexosamine-containing polysaccharide intracellular adhesion. *Infect. Immun.* **62**:3244–3253.
- Maira-Litran, T., A. Kropec, C. Abeygunawardana, J. Joyce, G. Mark III, D. A. Goldmann, and G. B. Pier. 2002. Immunochemical properties of the staphylococcal poly-N-acetylglucosamine surface polysaccharide. *Infect. Immun.* **70**:4433–4440.
- Marrie, T. J., and J. W. Costerton. 1984. Scanning and transmission electron microscopy of in situ bacterial colonization of intravenous and intraarterial catheters. *J. Clin. Microbiol.* **19**:687–693.
- Marrie, T. J., J. Nelligan, and J. W. Costerton. 1982. A scanning and transmission electron microscopic study of an infected endocardial pacemaker lead. *Circulation* **66**:1339–1341.
- Novick, R. 1967. Properties of a cryptic high-frequency transducing phage in *Staphylococcus aureus*. *Virology* **33**:155–166.
- O'Reilly, M., J. C. de Azavedo, S. Kennedy, and T. J. Foster. 1986. Inactivation of the alpha-haemolysin gene of *Staphylococcus aureus* 8325–4 by site-directed mutagenesis and studies on the expression of its haemolysins. *Microb. Pathog.* **1**:125–138.
- Song, L., M. R. Hobaugh, C. Shustak, S. Cheley, H. Bayley, and J. E. Gouaux. 1996. Structure of staphylococcal alpha-hemolysin, a heptameric transmembrane pore. *Science* **274**:1859–1866.
- Vuong, C., H. L. Saenz, F. Gotz, and M. Otto. 2000. Impact of the *agr* quorum-sensing system on adherence to polystyrene in *Staphylococcus aureus*. *J. Infect. Dis.* **182**:1688–1693.
- Walker, B., M. Krishnasastri, L. Zorn, and H. Bayley. 1992. Assembly of the oligomeric membrane pore formed by Staphylococcal alpha-hemolysin examined by truncation mutagenesis. *J. Biol. Chem.* **267**:21782–21786.