

Dartmouth College

## Dartmouth Digital Commons

---

Open Dartmouth: Published works by  
Dartmouth faculty

Faculty Work

---

5-17-2005

### Cellulose Utilization by Clostridium Thermocellum: Bioenergetics and Hydrolysis Product Assimilation

Yi-Heng P. Zhang  
*Dartmouth College*

Lee R. Lynd  
*Dartmouth College*

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Biological Engineering Commons](#), [Biology Commons](#), and the [Environmental Microbiology and Microbial Ecology Commons](#)

---

#### Dartmouth Digital Commons Citation

Zhang, Yi-Heng P. and Lynd, Lee R., "Cellulose Utilization by Clostridium Thermocellum: Bioenergetics and Hydrolysis Product Assimilation" (2005). *Open Dartmouth: Published works by Dartmouth faculty*. 1382. <https://digitalcommons.dartmouth.edu/facoa/1382>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Published works by Dartmouth faculty by an authorized administrator of Dartmouth Digital Commons. For more information, please contact [dartmouthdigitalcommons@groups.dartmouth.edu](mailto:dartmouthdigitalcommons@groups.dartmouth.edu).

# Cellulose utilization by *Clostridium thermocellum*: Bioenergetics and hydrolysis product assimilation

Yi-Heng Percival Zhang\* and Lee R. Lynd\*<sup>††</sup>

\*Thayer School of Engineering and <sup>†</sup>Department of Biological Sciences, Dartmouth College, Hanover, NH 03755

Edited by Arnold L. Demain, Drew University, Madison, NJ, and approved April 1, 2005 (received for review November 23, 2004)

The bioenergetics of cellulose utilization by *Clostridium thermocellum* was investigated. Cell yield and maintenance parameters,  $Y_{X/ATP}^{True} = 16.44$  g cell/mol ATP and  $m = 3.27$  mmol ATP/g cell per hour, were obtained from cellobiose-grown chemostats, and it was shown that one ATP is required per glucan transported. Experimentally determined values for  $G_{ATP}^{P-T}$  (ATP from phosphorylytic  $\beta$ -glucan cleavage minus ATP for substrate transport, mol ATP/mol hexose) from chemostats fed  $\beta$ -glucans with degree of polymerization (DP) 2–6 agreed well with the predicted value of  $(n-1)/n$  ( $n = \text{mean cellobiose DP assimilated}$ ). A mean  $G_{ATP}^{P-T}$  value of  $0.52 \pm 0.06$  was calculated for cellulose-grown chemostat cultures, corresponding to  $n = 4.20 \pm 0.46$ . Determination of intracellular  $\beta$ -glucan radioactivity resulting from  $^{14}\text{C}$ -labeled substrates showed that uptake is different for cellulose and cellobiose (G2). For  $^{14}\text{C}$ -cellobiose, radioactivity was greatest for G2; substantially smaller but measurable for G1, G3, and G4; undetectable for G5 and G6; and  $n$  was  $\approx 2$ . For  $^{14}\text{C}$ -cellulose, radioactivity was greatest for G5; lower but substantial for G6, G2, and G1; very low for G3 and G4; and  $n$  was  $\approx 4$ . These results indicate that: (i) *C. thermocellum* hydrolyzes cellulose by a different mode of action from the classical mechanism involving solubilization by cellobiohydrolase; (ii) bioenergetic benefits specific to growth on cellulose are realized, resulting from the efficiency of oligosaccharide uptake combined with intracellular phosphorylytic cleavage of  $\beta$ -glucosidic bonds; and (iii) these benefits exceed the bioenergetic cost of cellulase synthesis, supporting the feasibility of anaerobic biotechnological processing of cellulosic biomass without added saccharolytic enzymes.

cellulose hydrolysis | cellulase | cellulosome | anaerobic | thermophilic | ABC transport

Biologically mediated cellulose hydrolysis is a major flow of carbon in the biosphere (1, 2), important in several agricultural processes (3), and could be widely used to produce sustainable fuels and chemicals (3–7). Extensive evidence obtained with noncomplexed cellulase systems produced by aerobic microorganisms supports a mechanism involving synergistic action by endoglucanases and cellobiohydrolases with cellobiose the main product of cellulose solubilization (8–10). Anaerobic microorganisms possess complexed cellulase systems with architecture distinct from the noncomplexed systems of aerobes (11–13). In cell-free experiments involving complexed cellulase systems, cellobiose and glucose have been observed to accumulate (14, 15), consistent with a hydrolysis mechanism similar to that for noncomplexed systems. No evidence has been presented that the primary product of cellulose hydrolysis utilized by anaerobic microorganisms is other than cellobiose.

Utilization of cellulose by anaerobic microorganisms is a challenging proposition from a bioenergetic perspective, because the modest ATP available from anaerobic catabolism needs to support both microbial growth and cellulase production (16). Because reaction rates catalyzed by cellulases are lower than for most other enzymes, typically by at least 2 orders of magnitude on a protein-specific basis (9, 10), large amounts of cellulase and hence ATP are required to achieve significant cellulose hydrolysis rates (17, 18). Existing bioenergetic models

of microbial cellulose utilization are not sufficient to definitively evaluate the extent of the “metabolic burden” associated with cellulase synthesis for either naturally occurring cellulolytic microorganisms or biotechnological processes.

Were such an evaluation available, it would provide fundamental insights into the physiology of anaerobic cellulolytic microorganisms, the ecological communities in which these organisms occur, and the evolutionary strategies they embody. It would also address a central factor determining the feasibility of industrial processes in which anaerobic microorganisms convert cellulosic biomass to a desired product in the absence of added saccharolytic enzymes. Such “consolidated bioprocessing” is a potential breakthrough for low-cost production of ethanol or other fermentation products from cellulosic biomass (16, 19, 20).

In this study, we validate a comprehensive bioenergetic model of cellulose utilization by *Clostridium thermocellum*, a thermophilic cellulolytic anaerobic bacterium that exhibits one of the highest growth rates on crystalline cellulose among described microorganisms (16). In so doing, insights are gained with respect to substrate assimilation during cellulose utilization, the mechanism of cellulose hydrolysis, evaluation of the bioenergetic cost associated with growth on cellulose as compared with soluble substrates, and the feasibility of consolidated bioprocessing.

## Materials and Methods

**Chemicals and Microbial Cultures.** All chemicals were reagent grade and were obtained from Sigma unless indicated otherwise. *C. thermocellum* ATCC 27405 has been maintained in our lab since 1983, as described (21–23). Chemically defined, MTC medium was prepared by combining six sterile solutions under a nitrogen atmosphere, as described (21–23). Cellodextrins were prepared by mixed acid hydrolysis and separated chromatographically, as reported (24).

**Continuous Fermentation.** Continuous fermentations on cellobiose ( $\approx 5$  g/liter) and microcrystalline cellulose (Avicel PH 105, FMC, Philadelphia) Avicel ( $\approx 5$  g/liter) were carried out in 2.5-liter round-bottom reactors (Applikon Dependable Instruments, Foster City, CA) with agitation via a marine impeller at 300 rpm. Continuous fermentation on soluble cellodextrins with degree of polymerization from 2 to 6 was also carried out in a 60-ml working volume jacketed glass fermentor (NDS Glass, Vineland, NJ), as described (23). Cultures were considered to be at steady state when the samples exhibited  $<5\%$  variation and no consistent increasing or decreasing trend over time. Each reported steady-state value is based on the average of at least four data points.

**Analysis of Fermentation Products.** The mass concentrations of cellulase and cells growing on Avicel and cellodextrins were calculated based on an indirect ELISA by using antibody raised

This paper was submitted directly (Track II) to the PNAS office.

<sup>††</sup>To whom correspondence should be addressed. E-mail: lee.lynd@dartmouth.edu.

© 2005 by The National Academy of Sciences of the USA

against a sequence from the *C. thermocellum* scaffoldin protein, as described (21, 23). Noncellulase extracellular protein is defined as supernatant protein concentration minus the supernatant cellulase concentration as determined by ELISA. The dry weight, cellulose conversion, and fermentation product concentration of culture samples were measured as described (21, 23).

**Source of  $^{14}\text{C}$ -Labeled Substrates.** Radiolabeled cellulose (50  $\mu\text{Ci}$ ; 1 Ci = 37 GBq), purchased from American Radiolabeled Chemicals (St. Louis), was extracted by using 90% ethanol followed by boiling, dilution using 40 mg of Avicel PH105, and then treated with 83% phosphoric acid to obtain a radiolabeled amorphous cellulose suspension (21). Radiolabeled cellobiose was prepared by using cellobiose phosphorylase in conjunction with radiolabeled glucose (Sigma) and glucose-1-phosphate, as described (25, 26). Labeled cellobiose was separated by TLC (Whatman LK6DF), as described (26, 27), and the specific activity of cellobiose was calibrated by the modified BCA method (28). Silica gel from TLC plates was scraped at a location indicated by unlabeled cellobiose controls, added to distilled water, incubated at 80°C for 30 min with repeated vortex mixing, and removed by centrifugation.

**Uptake of  $^{14}\text{C}$ -Labeled Substrates.** Prewarmed oxygen-free labeled substrate (either cellulose or cellobiose) prepared as above was injected into 100-ml serum vials containing 50 ml of Avicel-grown cell cultures in late stationary phase (substrate recently exhausted) as well as a cell-free control containing *C. thermocellum* cellulase purified by affinity digestion (21, 29) at the same concentration present in the supernatant of the Avicel-grown culture. Samples were withdrawn at indicated intervals by using an 18-gauge needle connected to a 60-ml syringe containing 40 ml of ultracold ( $-70^\circ\text{C}$ ) methanol-Hepes buffer to quench metabolism immediately (30). Intracellular cellobioses were extracted by boiling in hot ethanol (30) and then treated by using 13% perchloric acid (final concentration) at 40°C for 1 h to convert glucose phosphate to glucose. The acidified metabolite solution was neutralized by a solution containing 2 M  $\text{K}_2\text{CO}_3$  and 2 M KOH (26). The concentrated metabolite extracts were applied to a TLC plate and run twice by using a developing solution, as described (26). The separated radiolabeled cellobioses were scraped from the TLC plate according to the locations of cold cellobiose standards (26, 27), and the radioactivities of individual cellobioses were measured by using a Beckman LS7500 liquid scintillation counter. The radioactivity of intracellular cellobioses was calculated by a procedure involving subtraction of noncell-associated cellulose-adhered labeled cellobioses inferred from a cell-free control experiment (see *Supporting Text*, which is published on the PNAS web site, for details).

## Results

ATP-producing metabolic processes available to *C. thermocellum* include intracellular phosphorolytic cleavage of  $\beta$ -glucosidic bonds by cellobioses and cellobiose phosphorylases (16, 21), glycolysis via the Emden-Meyerhoff pathway, and the action of acetate kinase. ATP-consuming metabolic processes include substrate transport via an adenosine-binding cassette system (16, 31), cell synthesis, cellulase synthesis, and nonbiosynthetic “maintenance” functions. For coupled ATP-limited metabolism, the rate of ATP production can be set equal to the rate of ATP consumption, resulting in Eq. 1 (ref. 16; see *Justification and explanation for Eq. 1* in *Supporting Text* for additional details):

$$r_S^X f \frac{n-1}{n} + r_S^X (Y_{E/\text{Glu}} + 2Y_{Ac/\text{Glu}} + Y_{L/\text{Glu}}) = r_S^X \frac{\alpha}{n} + \frac{D}{Y_{X/\text{ATP}}^{\text{True}}} \left( 1 + \frac{Y_{E/X}}{R} \right) + m, \quad [1]$$




Fig. 1. Plot for determination of  $Y_{X/\text{ATP}}^{\text{True}}$  and  $m$  based on Eq. 2. Data are calculated from steady-state continuous cultures of *C. thermocellum* growing on cellobiose with a feed concentration of  $\approx 5$  g/liter (see Table 3 which is published as supporting information on the PNAS web site, for details).

with  $f$  the fraction of  $\beta$ -glucosidic bonds cleaved phosphorolytically;  $n$ , mean degree of polymerization of assimilated cellobioses;  $Y_{E/\text{Glu}}$ ,  $Y_{Ac/\text{Glu}}$ , and  $Y_{L/\text{Glu}}$ , molar fermentation product yields (ethanol, acetate, and lactate, mol product/mol glucose);  $\alpha$ , ATP expenditure/cellobioses transported;  $r_S^X$ , cell-specific carbohydrate uptake rate (mol glucose equivalent per g cell per h);  $D$ , the dilution rate (1/h);  $Y_{E/X}$ , cell-specific yield of cellulase and additional supernatant protein (g protein/g cell);  $Y_{X/\text{ATP}}^{\text{True}}$ , true cell yield ATP (g cell/mol ATP used for anabolism);  $R$ , ratio of the true protein yield to the true cell yield; and  $m$ , maintenance (mol ATP/g cell per h). The literature supports a value of  $R = 0.82$  for bacterial growth (32, 33), and  $f = 1$  for *C. thermocellum* (22). We conclude that  $\alpha = 1$  for cellobioses and cellobioses transport by *C. thermocellum* based on statistical analysis of  $Y_{X/\text{ATP}}^{\text{True}}$  values (Table 2, which is published as supporting information on the PNAS web site). Several studies of adenosine-binding cassette transport systems under energy-limited conditions have also found that  $\alpha = 1$  (34–36).

For cellobioses ( $n = 2$ ) and substituting values for  $\alpha$ ,  $R$ , and  $f$ , Eq. 1 can be rewritten as


$$q_{\text{ATP, Cellobiose}} = r_S^X (Y_{E/\text{Glu}} + 2Y_{Ac/\text{Glu}} + Y_{L/\text{Glu}}) = \frac{D}{Y_{X/\text{ATP}}^{\text{True}}} \left( 1 + \frac{Y_{E/X}}{0.82} \right) + m, \quad [2]$$

where  $q_{\text{ATP, Cellobiose}}$  denotes the net rate of ATP synthesis. A plot of  $q_{\text{ATP, Cellobiose}}$  vs. the biosynthesis rate from cellobioses chemostat data exhibits a linear trend (Fig. 1), consistent with Eq. 2. From the slope and intercept of this line,  $Y_{X/\text{ATP}}^{\text{True}} = 16.44$  g cell/mol ATP and  $m = 3.27$  mmol/g cell per h.

For cellulose, with the mean cellobioses chain length assimilated,  $n$ , unknown, Eq. 1 can be rewritten as

$$G_{\text{ATP}}^{P-T} = \frac{n-2}{n} = \frac{D}{r_S^X Y_{X/\text{ATP}}^{\text{True}}} \left( 1 + \frac{Y_{E/X}}{0.82} \right) + \frac{m}{r_S^X} - Y_{E/\text{Glu}} - 2Y_{Ac/\text{Glu}} - Y_{L/\text{Glu}}, \quad [3]$$

where  $G_{\text{ATP}}^{P-T}$  corresponds to the ATP gained by phosphorolytic cleavage minus ATP expended on substrate transport (mol ATP/mol hexose). A plot of  $G_{\text{ATP}}^{P-T}$  vs.  $n$  (Fig. 2) shows excellent agreement between the model prediction,  $G_{\text{ATP}}^{P-T} = (n-2)/n$ , and experimental values calculated from steady-state chemostat cultures grown on purified cellobioses of length 2–6 at  $D =$




**Fig. 2.** Predicted (○) and measured  $G_{ATP}^{P-T}$  (●) values as a function of  $n$ . Experimental points are calculated from steady-state continuous cultures of *C. thermocellum* growing on cellodextrins of length 2–6 as well as cellulose (Avicel). The dashed line denotes the mean value of  $G_{ATP}^{P-T}$  for cellulose fermentation, with the shaded region denoting the standard deviation for independent steady-state fermentation runs carried out at various dilution rates (see Tables 4 and 5, which are published as supporting information on the PNAS web site, for details).

$0.167 \text{ h}^{-1}$ . For growth on cellulose, an average  $G_{ATP}^{P-T}$  value of  $0.52 \pm 0.06$  was calculated for chemostat cultures with  $D$  from  $0.0333$  to  $0.167 \text{ h}^{-1}$ . Solving Eq. 3 for  $n$  gives  $n = 4.20 \pm 0.46$ .

If the mean chain length of cellulose hydrolysis products taken up by *C. thermocellum* is substantially greater than 2, as indicated by the results presented thus far, then one would expect to see significantly different distributions of labeled intracellular cellodextrins accompanying growth on  $^{14}\text{C}$ -cellulose as compared with  $^{14}\text{C}$ -labeled cellobiose. On the other hand, if cellobiose were the primary hydrolysis product, as per the classical mechanism, then little or no difference between cellobiose- and cellulose-labeled substrates should be observed. With these observations in mind,  $^{14}\text{C}$ -labeled substrates, either cellobiose or cellulose, were added to substrate-exhausted cellulose-grown cultures, and the radioactivity of intracellular labeled cellodextrins was monitored. For labeled cellobiose added (Fig. 3A), the greatest amount of radioactivity is measured for cellobiose (G2) with lesser amounts for G1, G3, and G4. Radioactive G5 and G6 are not detected, and the mean cellodextrin degree of polymerization (DP) is  $\approx 2$ . For labeled cellulose, the greatest amount of radioactivity is detected for G5, with somewhat lesser amounts for G6, G2, and G1, and the lowest amounts for G3 and G4. The mean cellodextrin DP is  $\approx 4$ . Although the cellodextrin pool sizes measured cannot be equated to fluxes through these pools, the data indicate that G5 and G6 are assimilated in significant amounts during growth on cellulose, whereas this is not the case for growth on cellobiose. The results in Fig. 3 provide independent confirmation of our prediction based on bioenergetic analysis that  $n$  is substantially greater than 2 during growth on cellulose. The appearance of G3 and G4 during growth on cellobiose is not unexpected, because cellobiose and cellodextrin phosphorylases are readily reversible (20).

Fig. 4 presents the rates of ATP generation (Fig. 4A) and consumption (Fig. 4B) as a function of growth rate for *C. thermocellum* grown on cellulose. These data are calculated by using  $Y_{X/ATP}^{\text{True}}$  and  $m$  determined from Fig. 1 with  $n = 4.2$ . Glycolysis is responsible for slightly over half of the ATP supply with the remainder split roughly equally between phosphorolysis and acetate kinase. Cell synthesis accounts for an increasing share of ATP consumption with increasing growth rate up to a




**Fig. 3.** Radioactivity of intracellular cellodextrins following addition of [ $^{14}\text{C}$ ]cellobiose ( $0.05 \mu\text{Ci/ml}$  broth, A) and phosphoric acid-swollen [ $^{14}\text{C}$ ]cellulose ( $0.5 \mu\text{Ci/ml}$  broth, B). ▲, glucose; ●, cellobiose; ■, cellotriose; ▼, cellotetraose; ●, cellopentaose; ■, cellohexaose; and ◆, number average degree of polymerization of intracellular cellodextrin.

maximum of 45%. ATP demands for cellulase synthesis and transport are relatively small ( $\leq 17\%$ ). Because significant quantities of noncellulase protein were measured in culture supernatants, perhaps due to cell lysis, corresponding ATP consumption is shown in Fig. 4B.

## Discussion

We conclude that *C. thermocellum* assimilates cellodextrins with  $n \approx 4$  during growth on cellulose, implying that the immediate products of cellulose solubilization have an average degree of polymerization of at least this value. The mechanism of cellulose hydrolysis by *C. thermocellum* thus is different from the accepted mechanism for aerobic microorganisms such as *Trichoderma reesei*, in which cellobiose is the primary direct product of cellulose hydrolysis. Although this result contradicts current understanding of cellulose utilization by *C. thermocellum*, as well as other cellulolytic microbes, it is consistent with a number of observations from prior studies. *C. thermocellum* is known to have an uptake system capable of assimilating cellodextrins from G2 to G6 (26). Of the 16 catalytically active cellulosome components whose mode of action has been determined (11–13), 12 are endoglucanases, and only 2 are strict cellobiohydrolases. Prior data do not contradict the hypothesis that the initial products of cellulose hydrolysis produced by the *C. thermocellum*





**Fig. 4.** ATP generation (A) and demand (B) for *C. thermocellum* growing on cellulose (Avicel). Data are calculated from steady-state continuous cultures of *C. thermocellum* growing on Avicel (see Table 5 for details).

cellulase system are higher molecular weight cellodextrins that are rapidly cleaved in the absence of cells and thus do not appear in significant concentrations in cell-free experiments but are available for microbial uptake (16). Recent results involving rapid sampling of several purified cellulase components support this interpretation (ref. 37; W. Schwartz, personal communication). The distance between two adjacent catalytic subunits in the *C. thermocellum* cellulosome has been estimated at eight glucosidic bonds (38). Thus, simultaneous catalytic events mediated by adjacent catalytic components would be expected to result in an insoluble G8 fragment, and any subsequent cleavage of this G8 fragment would result in two soluble products with mean chain length 4.

*C. thermocellum* utilizes a relatively narrow range of substrates and appears to have evolved to function as a cellulose-using “specialist” (16, 31). Cellodextrin transport using an adenosine-binding cassette system, which typically features very high affinity [e.g.,  $K_m = 52 \mu\text{M}$  for cellobiose (26)], puts the organism in a strong position to compete for products of cellulose hydrolysis and is responsive to demands associated with growth under high-temperature and nutrient-limited conditions (39, 40). By assimilating cellodextrins with length  $\approx 4$  during growth on cellulose, the organism avoids an otherwise rather high ATP requirement for an anaerobe to expend on substrate transport. An equally large additional benefit of such assimilation is

**Table 1.** Net ATP availability associated with metabolic processes impacted by substrate chain length

Process	ATP/mol glucose	
	Cellobiose	Cellulose
Phosphorolytic cleavage	+0.5	+0.75
Transport	-0.5	-0.25
Cellulase synthesis	-0.1 to -0.2	-0.26 to -0.32
Net	-0.1 to -0.2	+0.18 to +0.24

Values for cellulose are based on  $n = 4.2$ , with ATP requirements for cellulase synthesis based on the range of values observed for cellulose-grown chemostats with  $D = 0.033$  to  $0.167 \text{ h}^{-1}$ .

realized by phosphorolytic cleavage of  $\beta$ -glucan bonds. *C. thermocellum* does not grow easily or well on glucose (16) and uses cellobiose instead of glucose if presented with both substrates (16, 31). This can readily be understood in light of the fact that no benefit is derived from phosphorolytic cleavage during growth on glucose, whereas the cost of transport per hexose, by either an adenosine-binding cassette system or any other active mechanism, is higher for glucose than for oligosaccharides. As shown in Table 1, the benefits of efficient transport and phosphorolytic cleavage of cellodextrins more than compensate for the higher ATP expended on cellulase synthesis when the organism is grown on cellulose as compared with cellobiose. In addition to cellobiose and cellodextrin phosphorylases, found in a variety of cellulolytic microorganisms as well as *C. thermocellum* (22), several other intracellular phosphorylases have been reported to cleave soluble oligosaccharides such as maltose and maltodextrins in the presence of inorganic phosphate to glucose-1-phosphate and  $G_{n-1}$  (35, 41, 42). Although this suggests that substrate level phosphorylation linked to phosphorolytic glucosidic bond cleavage may be widespread, the quantitative significance of these enzyme activities remains to be evaluated in the context of overall ATP supply in the organisms that produce them.

From a biotechnological perspective, our results establish that consolidated bioprocessing of cellulose is bioenergetically feasible for fermentative production of ethanol or other products that do not require ATP for synthesis from intermediates of central metabolism. The strategy of assimilating oligosaccharides combined with phosphorolytic cleavage may be of value for biotechnological application beyond cellulose conversion. Engineering these features into microbial biocatalysts and using widely available feedstocks containing soluble oligosaccharides would provide extra ATP for microbial growth, product synthesis, and/or transport. Although such extra ATP represents a small fraction of the total available from aerobic respiration, it could make a significant and potentially enabling difference for applications involving high yields of reduced products and for processes based on anaerobic fermentation in particular.

We thank P. J. Weimer, W. Schwarz, and H. J. Strobel for useful discussions, and we are grateful to Michael Campbell for editorial support. We also acknowledge support from the Department of Energy (DE-FG02-02ER15350) and the National Institute of Standards and Technology (60NANBID0064).

- Berner, R. A. (2003) *Nature* **426**, 323–326.
- Melillo, J. M., Stuedler, P. A., Aber, J. D., Newkirk, K., Lux, H., Bowles, F. P., Catricala, C., Magill, A., Ahrens, T. & Morrisseau, S. (2002) *Science* **298**, 2173–2176.
- Russell, J. B. & Rychlik, J. L. (2001) *Science* **292**, 1119–1122.
- Kamm, B. & Kamm, M. (2004) *Appl. Microbiol. Biotechnol.* **64**, 137–145.
- Wahlund, B., Yan, J. & Westermarck, M. (2004) *Biomass Bioenerg.* **26**, 531–544.
- Wirth, T. E., Gray, C. B. & Podesta, J. D. (2003) *Foreign Aff.* **82**, 125–144.

- Wyman, C. E. (2003) *Biotechnol. Prog.* **19**, 254–262.
- Divne, C., Stahlberg, J., Reinikainen, T., Ruohonen, L., Pettersson, G., Knowles, J. K., Teeri, T. T. & Jones, T. A. (1994) *Science* **265**, 524–528.
- Teeri, T. T., Koivula, A., Linder, M., Wohlfahrt, G., Divne, C. & Jones, T. A. (1998) *Biochem. Soc. Trans.* **26**, 173–178.
- Zhang, Y.-H. P. & Lynd, L. R. (2004) *Biotechnol. Bioeng.* **88**, 797–824.
- Bayer, E. A., Belaich, J.-P., Shoham, Y. & Lamed, R. (2004) *Annu. Rev. Microbiol.* **58**, 521–554.

12. Doi, R. H. & Kosugi, A. (2004) *Nat. Rev. Microbiol.* **2**, 541–551.
13. Schwarz, W. H. (2001) *Appl. Microbiol. Biotechnol.* **56**, 634–649.
14. Johnson, E. A., Reese, E. T. & Demain, A. L. (1982) *J. Appl. Biochem.* **4**, 64–71.
15. Hom-nami, K., Coughlan, M. P., Hon-nami, H. & Ljungdahl, L. G. (1986) *Arch. Microbiol.* **145**, 13–19.
16. Lynd, L. R., Weimer, P. J., van Zyl, W. H. & Pretorius, I. S. (2002) *Microbiol. Mol. Biol. Rev.* **66**, 506–577.
17. Zhou, S. D., Davis, F. C. & Ingram, L. O. (2001) *Appl. Environ. Microbiol.* **67**, 6–14.
18. van Walsum, G. P. & Lynd, L. R. (1998) *Biotechnol. Bioeng.* **58**, 316–320.
19. Office of the Biomass Program of U.S. Department of Energy (2003) *Biomass Program: Multi-Year Technical Plan FY04–FY08*, [www.nrel.gov/docs/fy05osti/36999.pdf](http://www.nrel.gov/docs/fy05osti/36999.pdf).
20. National Research Council (1999) *Review of the Research Strategy for Biomass-Derived Transportation Fuels* (Natl. Acad. Press, Washington, DC).
21. Zhang, Y. & Lynd, L. R. (2003) *Anal. Chem.* **75**, 219–227.
22. Zhang, Y.-H. P. & Lynd, L. R. (2004) *Appl. Environ. Microbiol.* **70**, 1563–1569.
23. Zhang, Y.-H. P. & Lynd, L. R. (2005) *J. Bacteriol.* **188**, 99–106.
24. Zhang, Y.-H. P. & Lynd, L. R. (2003) *Anal. Biochem.* **322**, 225–332.
25. Ng, T. K. & Zeikus, J. G. (1986) *Appl. Environ. Microbiol.* **52**, 902–905.
26. Strobel, H. J., Caldwell, F. C. & Dawson, K. A. (1995) *Appl. Environ. Microbiol.* **61**, 4012–4015.
27. Chirico, W. J. & Brown, R. D., Jr. (1985) *Anal. Biochem.* **150**, 264–272.
28. Zhang, Y.-H. P. & Lynd, L. R. (2005) *Biomacromolecules* **6**, in press.
29. Morag, E., Bayer, E. A. & Lamed, R. (1992) *Enzyme Microb. Technol.* **14**, 289–295.
30. Raamsdonk, L. M., Teusink, B., Broadhurst, D., Zhang, N., Hayes, A., Walsh, M. C., Berden, J. A., Brindle, K. M., Kell, D. B., Rowland, J. J., *et al.* (2001) *Nat. Biotechnol.* **19**, 45–50.
31. Mitchell, W. J. (1998) *Adv. Microb. Physiol.* **39**, 31–130.
32. Forrest, W. W. & Walker, D. J. (1971) *Adv. Microb. Physiol.* **5**, 213–274.
33. Russell, J. B. & Cook, G. M. (1995) *Microbiol. Rev.* **9**, 48–62.
34. Mimmack, M. L., Gallagher, M. P., Pearce, S. R., Hyde, S. C., Booth, I. R. & Higgins, C. F. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 8257–8261.
35. Muir, M., Williams, L. & Ferenci, T. (1985) *J. Bacteriol.* **163**, 1237–1242.
36. Shapiro, A. B. & Ling, V. (1998) *Eur. J. Biochem.* **254**, 189–193.
37. Zverlov, V. V., Velikodvorskaya, G. A. & Schwarz, W. H. (2003) *Microbiology* **149**, 515–524.
38. Mayer, F., Coughlan, M. P., Mori, Y. & Ljungdahl, L. G. (1987) *Appl. Environ. Microbiol.* **53**, 2785–2792.
39. Albers, S. V., Van de Vossen, J. L., Driessen, A. J. & Konings, W. N. (2001) *Extremophiles* **5**, 285–294.
40. Ferenci, T. (1999) *Curr. Opin. Microbiol.* **2**, 208–213.
41. Schinzel, R. & Nidetzky, B. (1999) *FEMS Microbiol. Lett.* **171**, 73–79.
42. Geremia, S., Campagnolo, M., Schinzel, R. & Johnson, L. N. (2002) *J. Mol. Biol.* **322**, 413–423.