

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

7-18-2006

A Subset of Arabidopsis AP2 Transcription Factors Mediates Cytokinin Responses in Concert with a Two-Component Pathway

Aaron M. Rashotte

University of North Carolina at Chapel Hill

Michael G. Mason

Dartmouth College

Claire E. Hutchison

University of North Carolina at Chapel Hill

Fernando J. Ferreira

University of North Carolina at Chapel Hill

G. Eric Schaller

Dartmouth College

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Biology Commons](#), [Genetics Commons](#), [Physiology Commons](#), and the [Plant Sciences Commons](#)

Dartmouth Digital Commons Citation

Rashotte, Aaron M.; Mason, Michael G.; Hutchison, Claire E.; Ferreira, Fernando J.; Schaller, G. Eric; and Kieber, Joseph J., "A Subset of Arabidopsis AP2 Transcription Factors Mediates Cytokinin Responses in Concert with a Two-Component Pathway" (2006). *Dartmouth Scholarship*. 1403.

<https://digitalcommons.dartmouth.edu/facoa/1403>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Authors

Aaron M. Rashotte, Michael G. Mason, Claire E. Hutchison, Fernando J. Ferreira, G. Eric Schaller, and Joseph J. Kieber

A subset of *Arabidopsis* AP2 transcription factors mediates cytokinin responses in concert with a two-component pathway

Aaron M. Rashotte*, Michael G. Mason†, Claire E. Hutchison*, Fernando J. Ferreira*, G. Eric Schaller†, and Joseph J. Kieber**

*Department of Biology, University of North Carolina, Chapel Hill, NC 27599-3280; and †Department of Biological Sciences, Dartmouth College, Hanover, NH 03755

Edited by Maarten Koornneef, Wageningen University and Research Centre, Wageningen, Netherlands, and approved June 1, 2006 (received for review March 13, 2006)

The plant hormone cytokinin regulates numerous growth and developmental processes. A signal transduction pathway for cytokinin has been elucidated that is similar to bacterial two-component phosphorelays. In *Arabidopsis*, this pathway is comprised of receptors that are similar to sensor histidine kinases, histidine-containing phosphotransfer proteins, and response regulators (ARRs). There are two classes of response regulators, the type-A ARRs, which act as negative regulators of cytokinin responses, and the type-B ARRs, which are transcription factors that play a positive role in mediating cytokinin-regulated gene expression. Here we show that several closely related members of the *Arabidopsis* AP2 gene family of unknown function are transcriptionally up-regulated by cytokinin through this pathway, and we have designated these AP2 genes *CYTOKININ RESPONSE FACTORS* (CRFs). In addition to their transcriptional regulation by cytokinin, the CRF proteins rapidly accumulate in the nucleus in response to cytokinin, and this relocalization depends on the histidine kinases and the downstream histidine-containing phosphotransfer proteins, but is independent of the ARRs. Analysis of loss-of-function mutations reveals that the CRFs function redundantly to regulate the development of embryos, cotyledons, and leaves. Furthermore, the CRFs mediate a large fraction of the transcriptional response to cytokinin, affecting a set of cytokinin-responsive genes that largely overlaps with type-B ARR targets. These results indicate that the CRF proteins function in tandem with the type-B ARRs to mediate the initial cytokinin response. Thus, the evolutionarily ancient two-component system that is used by cytokinin branches to incorporate a unique family of plant-specific transcription factors.

cell signaling | plant hormones

Cytokinins are N^6 substituted adenine derivatives that were first identified by their ability to promote division in cultured plant cells together with a second hormone, auxin (1). Cytokinins have since been shown to play a role in diverse aspects of plant growth and development, including cell division, shoot initiation, apical meristem function, and vascular formation (2, 3). Recently, remarkable progress has been made in our understanding of cytokinin biosynthesis, metabolism, and perception. The genes encoding the key biosynthetic enzymes have been identified in plants, as have genes encoding several important cytokinin metabolic enzymes (4). A model for cytokinin perception and signaling has emerged that is similar to bacterial two-component phosphorelays (5). Binding of cytokinins to the *Arabidopsis* sensor histidine kinases (AHKs) initiates a phosphorelay in which the *Arabidopsis* histidine-containing phosphotransfer proteins (AHPs) are phosphorylated and then translocate into the nucleus where they likely transfer the phosphate to the *Arabidopsis* type-B response regulators (ARRs) (6–10). The type-B ARRs play a role in mediating the transcriptional response to cytokinin, including the induction of a second class of response regulators called the type-A ARRs (6, 11). The

type-A ARRs are cytokinin primary response genes that act as highly redundant negative regulators of the primary signal transduction pathway (12–14). The pseudophosphotransfer protein AHP6 also acts as a negative regulator of cytokinin signaling, and plays a role in vascular development (15).

Gene expression in response to cytokinin has been extensively studied, and numerous genes have been identified that are transcriptionally up-regulated in response to cytokinin (16–18) including two members of the AP2/ERF superfamily of transcription factors within the ethylene response factor (ERF) family (17). Here, we demonstrate that this subgroup of AP2 transcription factors moves into the nucleus in response to cytokinin, and that they mediate, together with the type-B ARRs, the transcriptional response to cytokinin.

Results and Discussion

The CRF Genes Are Transcriptionally Induced by Cytokinin in a Type-B ARR-Dependent Manner. The ERF family is comprised of 65 genes (19), several of which have been implicated in the response to the plant hormones ethylene, jasmonic acid, and cytokinin (20–22). Phylogenetic analysis of the predicted ERF-like proteins places the two cytokinin up-regulated genes in a subclade of six previously uncharacterized genes that fall into three pairs based on sequence similarity (Fig. 1*B*; see Figs. 5 and 6, which are published as supporting information on the PNAS web site). We named these six genes *CYTOKININ RESPONSE FACTORS* (CRFs) to reflect their similarity to ERFs and the observation that some members are transcriptionally up-regulated by cytokinin.

We examined the expression of the CRF genes in response to cytokinin by northern analysis. Consistent with previously reported microarray data (see Table 2, which is published as supporting information on the PNAS web site), the CRF2 and CRF5 transcripts are up-regulated 2- to 4-fold by cytokinin (Fig. 1*A*). The induction of both genes is rapid (<30 min), although CRF2 expression peaks at ≈60 min and then begins to decline, whereas induction of CRF5 is more sustained. CRF6 expression is also up-regulated by cytokinin, but more slowly. In contrast, the transcript levels of CRF1, CRF3, and CRF4 show little or no change in response to cytokinin (Fig. 1*A* and Table 2). The induction of CRF2 and CRF5 by cytokinin depends on the type-B ARRs, as an *arr1,12* double mutant (23) severely reduces the response of these genes to cytokinin (Fig. 1*C*). Examination of

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: AHK, *Arabidopsis* histidine kinase; AHP, *Arabidopsis* histidine-containing phosphotransfer protein; ARR, *Arabidopsis* response regulator; ERF, ethylene response factor; CRF, cytokinin response factor; CaMV, cauliflower mosaic virus.

†To whom correspondence should be addressed. E-mail: jkieber@unc.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Induction of *CRF* genes by cytokinin. (A) Levels of *CRF* transcripts after cytokinin treatment were examined by using Northern blot analysis. (B) An unrooted phylogenetic tree of the 65 ERF predicted proteins. Representative proteins are indicated; for a full list, see Fig. 5. The CRFs comprise a distinct subfamily with the cytokinin inducible members indicated by an asterisk. (C) Cytokinin induction of *CRF* transcripts requires type-B ARR. Real-time PCR of *CRF2* and *CRF5* confirm that they are induced after cytokinin treatment (10 μ M BA) for 1 h relative to a DMSO control in wild type, but are not induced in the type-B mutant *arr1,12*.

the 5' region of the *CRF2* and *CRF5* genes reveals the presence of multiple type-B consensus binding sites (17, 24), consistent with these CRFs being direct targets of type-B ARRs.

CRF Proteins Rapidly Relocalize to the Nucleus in Response to Cytokinin. To explore the subcellular localization of the CRF proteins, we examined protoplasts transformed with GFP:CRF fusions expressed from the cauliflower mosaic (CaMV) 35S promoter. In the absence of cytokinin, we observed relatively uniform fluorescence throughout the cell. After treatment with cytokinin, all six GFP:CRF fusion proteins accumulated in the nucleus (Fig. 2A; see Fig. 7, which is published as supporting information on the PNAS web site). This nuclear accumulation was never observed in protoplasts treated with a DMSO vehicle control. Nuclear accumulation in response to cytokinin was observed in >50% of the transformed protoplasts and was very rapid, occurring within 5–10 min, which is consistent with a relocation of preexisting protein. Additionally, this characteristic is independent of transcriptional induction, as these CRF fusions were driven by the constitutive CaMV 35S promoter. Cytokinin-induced nuclear accumulation is not a general property of ERF transcription factors, as two other ERFs that cluster outside of the CRF subgroup display constitutive nuclear localization (25). The rapid cytokinin-induced intracellular relocalization of CRFs was also observed in wild-type *Arabidopsis* stably transformed with *CRF2*:GFP, *CRF5*:GFP, or *CRF6*:GFP translation fusions expressed from their native promoters (Fig. 2C; data not shown). This relocalization was observed in the majority of the root hair cells that were examined, and occurred in multiple other cell types as well (data not shown). Consistent with the rapid kinetics, treatment with cycloheximide (an inhibitor of protein synthesis) did not inhibit nuclear accumulation of *CRF5*:GFP in seedlings (Fig. 2C) or in protoplasts (data not shown), indicating that preexisting CRF protein moves into the nucleus.

CRF Protein Relocalization Defines a Branch Point in the Cytokinin Phosphorelay Response Pathway. We examined the role of two-component elements in the cytokinin-induced nuclear accumulation of the CRFs. To this end, CaMV 35S:GFP:CRF2 constructs were transformed into protoplasts derived from cytokinin signaling mutants. In homozygous double *ahk3,4* and triple *ahk2,3,4* receptor mutant protoplasts, the distribution of the fluorescence signal from the GFP:CRF2 fusion protein was not altered by the addition of cytokinin (Fig. 2B), indicating that cytokinin-mediated nuclear accumulation of the CRFs requires the cytokinin AHK receptors. The histidine kinase inhibitor 3,3',4',5-tetrachlorosalicylanilide (TCSA) also blocked the movement of 35S:GFP:CRF2 in response to cytokinin (Fig. 2B), indicating that the histidine kinase activity of the cytokinin receptors is required for relocalization.

Functioning immediately downstream of the cytokinin receptors are the AHPs, which like the CRFs, also exhibit cytokinin-dependent nuclear localization (6). As with the receptor triple mutants, no nuclear accumulation of GFP:CRF2 was observed in response to cytokinin in transformed *ahp1,2,3,4,5* protoplasts (C.E.H. and J.J.K., unpublished data) (Fig. 2B). In contrast, disruption of multiple type-B ARRs (i.e., the *arr1,2,10,12* mutant) did not affect CRF nuclear movement in response to cytokinin (Fig. 2B). This is unlikely a result of residual cytokinin responsiveness as the *arr1,2,10,12* mutant is more insensitive to cytokinin in a variety of cytokinin response assays (data not shown) than is the *ahk3,4* double mutant, which blocks the CRF nuclear accumulation. Likewise, disruption of multiple type-A ARRs (*arr3,4,5,6,8,9*) did not alter the distribution of the CRF proteins (Fig. 2B). Thus, both the AHPs and cytokinin receptors are required for the cytokinin-regulated movement of CRFs into the nucleus, but the downstream ARRs are not.

cells vs. *crf1,2,5* 1.0 ± 0.1 mm² and 705 ± 96 cells in 10-day-old seedlings). A triple cytokinin receptor mutant (*ahk2,3,4*) also displays reduced cotyledon size that results primarily from reduced cell expansion (*ahk2,3,4* cotyledon size is 1.9 ± 0.2 mm² and are comprised of 933 ± 21 cells in 10-day-old seedlings). The lack of cell expansion in the triple receptor mutant cotyledons is distinct from previous studies that found that the reduced size of the seventh leaf of the *ahk2,3,4* mutants was the result of reduced cell number (28), which likely reflects distinct effects of the *ahk* mutations on these two different organs. However, the severely affected *crf* triple mutant cotyledons have additional defects not seen in the cytokinin receptor knockouts, such as the lack of normal pigmentation, suggesting that there are aspects of this phenotype that may be cytokinin independent.

Interestingly, cytokinins have been closely linked to cotyledon expansion, and this expansion was used as a bioassay to quantify cytokinin until the advent of MS-based methods. The CRFs may mediate the effect of cytokinin on cotyledon cell expansion. In addition to cotyledons, the *crf* mutants also occasionally affect juvenile leaves, but otherwise do not have any other obvious effect on morphology or development, except for the embryo-lethal phenotype of the *crf5, crf6* double mutant discussed below.

In contrast to other *crf* mutant combinations tested, all combinations of independent alleles of both *crf5* and *crf6* resulted in embryo lethality that was fully penetrant (Fig. 3B; data not shown). In self-fertilized *crf5-1/crf5-1 crf6-1/CRF6*, *crf5-2/crf5-2 crf6-1/CRF6*, or *crf5-1/CRF5 crf6-2/crf6-2* plants, $\approx 25\%$ of the seeds are absent from the silique (Fig. 3B). Approximately 25% of the embryos in these siliques never progress beyond the late globular to early heart stage of development (Fig. 3C). However, none of the single *crf5* and *crf6* mutant alleles display any detectable embryo defects. In addition, the *crf5-1/crf5-1 crf6-1/crf6-1* embryo lethal phenotype can be fully complemented by transformation with a genomic *CRF5* fragment (Fig. 3B). These results suggest that the *CRF5* and *CRF6* genes are redundant and necessary for embryo development.

Unlike mutants in the *Arabidopsis* two-component signaling pathway, we found little effect of *crf* single, double and triple mutants in other cytokinin response assays. For example, there is little or no significant effect of these mutants on the response of seedlings to cytokinin in root elongation (see Fig. 9, which is published as supporting information on the PNAS web site), and only very minor effects on *in vitro* shoot initiation assays (see Fig. 10, which is published as supporting information on the PNAS web site). Furthermore, the morphology of *crf* mutant seedlings grown on cytokinin did not differ substantial from the wild type (data not shown).

The CRFs Mediate Gene Expression in Response to Cytokinin Together with the Type-B ARRs. Because the CRFs are predicted to function in the regulation of gene expression, we examined the response of the transcriptome to cytokinin in the *crf* mutants. Wild-type




Fig. 4. CRFs act in parallel with type-B ARRs to mediate cytokinin regulated gene expression. (A) Wild-type, *arr1,12*, *crf1,2,5*, and *crf2,3,6* seedlings were treated with either 10 μ M BA or a DMSO control for 1 h and gene expression analyzed by using a microarray. Genes that displayed a ≥ 2 -fold change in response to cytokinin in the wild type are shown. (B) Venn diagram of the 135 cytokinin-regulated genes affected by the *arr1,12*, *crf1,2,5*, and/or *crf2,3,6* mutations. (C) Model of cytokinin signaling. Both AHPs and CRFs move into the nucleus in response to cytokinin. Once there, the AHPs phosphorylate the type-B ARRs, which, together with CRFs, mediate cytokinin-regulated gene expression. See text for further details.

and *crf* mutant seedlings were treated with cytokinin for 1 h and gene expression examined in triplicate by using a 29,000 element *Arabidopsis* oligonucleotide microarray (Fig. 4A; see Table 4 and Fig. 11, which are published as supporting information on the PNAS web site). Strikingly, of 169 genes whose transcript level was altered by cytokinin in wild-type seedlings, 93 (55%) exhibited a 2-fold or greater decrease in their responsiveness in the *crf1,2,5* mutant, and 81 (48%) in the *crf2,3,6* mutant (Fig. 4B). This indicates that *CRF* genes regulate a significant portion of the early transcriptional response to cytokinin. The effect of the *crf* triple mutants on several of these genes was verified by using quantitative PCR (Table 1).

Comparison of the cytokinin response in a type-B ARR double mutant and the *crf* mutants revealed substantial overlap among the genes regulated by these two divergent families of transcription factors. In the *arr1,12* mutant, 111 genes (66%) exhibited a 2-fold or greater decrease in their cytokinin responsiveness as compared to wild type, with 68% of those affected in *arr1,12* similarly affected in the *crf1,2,5* mutant and 60% in the *crf2,3,6* mutant (Fig. 4B). Thus, the *CRF* and type-B ARR genes regulate an overlapping set of cytokinin-response genes. How-

Table 1. Real-time PCR confirmation of cytokinin-regulated microarray results

Gene name	At number	Wild type + vs. - BA		<i>crf1,2,5</i> + vs. - BA		<i>crf2,3,6</i> + vs. - BA	
		Microarray	RT-PCR	Microarray	RT-PCR	Microarray	RT-PCR
Zinc Finger B-box	At1g68520	3.1	2.8 ± 0.3	1.0	1.3 ± 0.1	1.1	1.1 ± 0.1
NAM AtNAC6	At4g27410	2.7	2.7 ± 0.2	1.3	0.7 ± 0.1	1.4	1.0 ± 0.1
Speckled POZ	At3g48360	3.5	2.7 ± 0.3	1.1	0.8 ± 0.1	0.9	1.0 ± 0.5
Senescence Assoc. P.	At1g53885	3.9	2.7 ± 0.3	1.3	1.7 ± 0.1	2.0	1.5 ± 0.3
ARR5	At3g48100	7.3	14.3 ± 0.4	3.2	26.6 ± 1.1	5.0	25.4 ± 0.9
ARR7	At1g19050	3.6	9.5 ± 0.3	5.7	15.1 ± 0.6	5.5	12.1 ± 0.4

Four genes regulated by cytokinin in wild type, but not in *CRF* mutants, were examined by using real-time PCR, as well as two type-A ARR genes and show similar trends to the microarrays in Fig. 4A.

ever, not all of these genes are similarly regulated, most notably the type-A *ARRs*, which exhibit reduced responsiveness to cytokinin in the *arr1,12* mutant, still exhibit a wild type or greater level of induction in the *crf1,2,5* and *crf2,3,6* mutants (Table 1; see Table 4). The differential regulation of the type-A *ARRs* may contribute to the difference in cytokinin-dependent phenotypes observed in the *crf* and type-B *arr* mutants.

Model for Cytokinin Response Pathway. These data are consistent with a model for cytokinin function illustrated in Fig. 4C. Cytokinin binding to the AHKs initiates a phosphorelay that results in the phosphorylation of the AHPs. The phosphorylated AHPs move into the nucleus to activate the type-B *ARRs*, which then increase the transcription of a subset of the CRFs. In parallel, the phosphorylated AHPs induce CRF proteins to accumulate in the nucleus. Thus, the relocalization of the CRF proteins defines a branch point in the cytokinin two-component signal transduction pathway. The activated CRFs, together with the activated type-B *ARRs*, mediate cytokinin-regulated gene expression, affecting an overlapping set of gene targets. In addition to mediating a portion of the cytokinin response initiated at the AHK receptors, the CRFs may also play a role in other signaling or developmental pathways, based on the difference in phenotypes associated with the *crf* loss-of-function mutations compared to those of the cytokinin receptor mutants (29, 30). In particular, the cytokinin receptor triple mutant is a viable plant that is distinct from the embryo lethal phenotype of the *crf5,crf6* double mutant. Thus, CRFs are likely to receive input either from other cytokinin receptors, or from non-cytokinin-dependent signaling sources.

Materials and Methods

Plant Materials and Treatments. Seedlings were grown under standard growth conditions as described (13), except for microarray experiments which were grown as described (23). Mutant lines are described in detail (see *Supporting Text*, which is published as supporting information on the PNAS web site). Cytokinin treatments were 1 μ M *N*⁶-benzyladenine (BA) for various times for Northern analysis, 2 μ M BA for 10 min for protoplast experiments, and 10 μ M for 1 h in microarray and

real-time PCR experiments (14-day-old plants). Cycloheximide was used at 50 μ M as 1h pretreatments. Stable transformation of *Arabidopsis* was performed by floral dip method (31). *Arabidopsis* leaf mesophyll protoplasts were isolated and transformed by electroporation as previously described (32) with minor modifications as detailed (see *Supporting Text*). Plasmids used for transformations are described (see *Supporting Text*).

Microscopy. Epifluorescent microscopy with Hoechst dye 33342 (1 ng·ml⁻¹) and an UV source were used to observe cells and identify nuclei. A GFP filter that blocks chlorophyll fluorescence and Hoechst dye 33342 fluorescence was used to examine localization of GFP fusion proteins. Further details of the microscopy are presented in *Supporting Text*.

Microarray Analysis. Microarray analyses were conducted by using the 29,000-element *Arabidopsis* oligonucleotide microarrays (<http://ag.arizona.edu/microarray>). Genes in these analyses were considered cytokinin regulated if they were found to be significant by a Welch's *t* test with an adjusted Bonferroni correction after sample/column normalization (TIGR MeV v3.1 available at www.tm4.org) and had a mean fold change ≥ 2 or ≤ -2 . Further details of the microarray analysis are given in *Supporting Text*.

Real-Time PCR. Real-time PCR was carried out (see *Supporting Text*). At least two biological and two sample replicates were performed for each treatment. *Ct* values were generated by subtracting blanks and the baseline average over cycles 1–10 with a 10 \times SD over the cycle range for each sample. Samples with melt curves that did not have a single distinct peak were excluded from further analysis. Fold-change of samples was calculated after normalization to β -tubulin *Ct* values from each sample.

We thank members of the J.J.K. laboratory for helpful suggestions. Financial support was provided by a Department of Energy grant (to J.J.K.), a National Institutes of Health grant (to J.J.K. and A.M.R.), a National Science Foundation grant (to J.J.K. and G.E.S.), and a U.S. Department of Agriculture/National Research Initiative Competitive Grants Program grant (to G.E.S.).

- Miller, C. O., Skoog, F., Von Saltza, M. H. & Strong, F. (1955) *J. Am. Chem. Soc.* **77**, 1392–1393.
- Mok, D. W. & Mok, M. C. (2001) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **89**, 89–118.
- Davies, P. J. (2004) *Plant Hormones: Biosynthesis, Signal Transduction, Action!* (Kluwer Academic, Dordrecht, The Netherlands).
- Kakimoto, T. (2003) *J. Plant Res.* **116**, 233–239.
- Ferreira, F. J. & Kieber, J. J. (2005) *Curr. Opin. Plant Biol.* **8**, 518–525.
- Hwang, I. & Sheen, J. (2001) *Nature* **413**, 383–389.
- Ueguchi, C., Sato, S., Kato, T. & Tabata, S. (2001) *Plant Cell Physiol.* **42**, 751–755.
- Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., Tabata, S., Shinozaki, K. & Kakimoto, T. (2001) *Nature* **409**, 1060–1063.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., Yamashino, T. & Mizuno, T. (2001) *Plant Cell Physiol.* **41**, 1017–1023.
- Suzuki, T., Miwa, K., Ishikawa, K., Yamada, H., Aiba, H. & Mizuno, T. (2001) *Plant Cell Physiol.* **42**, 107–113.
- Sakai, H., Honma, T., Aoyama, T., Sato, S., Kato, T., Tabata, S. & Oka, A. (2001) *Science* **294**, 1519–1521.
- Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Kiba, T., Ueguchi, C., Sugiyama, T. & Mizuno, T. (1999) *Plant Cell Physiol.* **40**, 733–742.
- To, J. P. C., Haberer, G., Ferreira, F. J., Deruère, J., Mason, M. G., Schaller, G. E., Alonso, J. M., Ecker, J. R. & Kieber, J. J. (2004) *Plant Cell* **16**, 658–671.
- Brandstatter, I. & Kieber, J. J. (1998) *Plant Cell* **10**, 1009–1020.
- Mähönen, A. P., Bonke, M., Kauppinen, L., Riikonon, M., Benfey, P. & Helariutta, Y. (2000) *Genes Dev.* **14**, 2938–2943.
- Kiba, T., Naitou, T., Koizumi, N., Yamashino, T., Sakakibara, H. & Mizuno, T. (2005) *Plant Cell Physiol.* **46**, 339–355.
- Rashotte, A. M., Carson, S. D. B., To, J. P. C. & Kieber, J. J. (2003) *Plant Physiol.* **123**, 184–194.
- Brenner, W. G., Romanov, G. A., Kollmer, I., Burkle, L. & Schmölling, T. (2005) *Plant J.* **44**, 314–333.
- Nakano, T., Suzuki, K., Fujimura, T. & Shinshi, H. (2006) *Plant Physiol.* **140**, 411–432.
- Ohme-Takagi, M. & Shinshi, H. (1995) *Plant Cell* **7**, 173–182.
- McGrath, K. C., Dombrecht, B., Manners, J. M., Schen, P. M., Edgar, C. I., Maclean, D. J., Scheible, W.-R., Udvardi, M. K. & Kazan, K. (2005) *Plant Physiol.* **139**, 949–959.
- van der Fits, L. & Memelink, J. (2001) *Plant J.* **25**, 43–53.
- Mason, M. G., Mathews, D. E., Argyros, D. A., Maxwell, B. B., Kieber, J. J., Alonso, J. M., Ecker, J. R. & Schaller, G. E. (2005) *Plant Cell* **17**, 3007–3018.
- Sakai, H., Aoyama, T. & Oka, A. (2000) *Plant J.* **24**, 703–711.
- Koroleva, O. A., Tomlinson, M. L., Leader, D., Shaw, P. & Doonan, J. H. (2005) *Plant J.* **41**, 162–174.
- Qi, Y., Sun, Y., Xu, L., Xu, Y. & Huang, H. (2004) *Planta* **219**, 270–276.
- Vroemen, C. W., Mordhorst, A. P., Albrecht, C., Kwaktaal, M. A. C. J. & de Vries, S. C. (2003) *Plant Cell* **15**, 1563–1577.
- Riefler, M., Novak, O., Strnad, M. & Schmölling, T. (2006) *Plant Cell* **18**, 40–54.
- Higuchi, M., Pischke, M. S., Mahonen, A. P., Miyawaki, K., Hashimoto, Y., Seki, M., Kobayashi, M., Shinozaki, K., Kato, T., Tabata, S., et al. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 8821–8826.
- Nishimura, C., Ohashi, Y., Sato, S., Kato, T., Tabata, S. & Ueguchi, C. (2004) *Plant Cell* **16**, 1365–1377.
- Clough, S. J. & Bent, A. F. (1998) *Plant J.* **16**, 735–743.
- Sheen, J. (1991) *Plant Cell* **3**, 225–245.