

6-1998

Identification of a Family of Zinc Transporter Genes from Arabidopsis that Respond to Zinc Deficiency

Natasha Grotz
Dartmouth College

Tama Fox
Dartmouth College


Erin Connolly
Dartmouth College

Walter Park
University of Minnesota - Duluth

Mary Lou Guerinot
Dartmouth College

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>

 Part of the [Biology Commons](#), [Molecular, Genetic, and Biochemical Nutrition Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Grotz, Natasha; Tama Fox; Connolly, Erin; Park, Walter; Guerinot, Mary Lou; and Eide, David, "Identification of a Family of Zinc Transporter Genes from Arabidopsis that Respond to Zinc Deficiency" (1998). *Open Dartmouth: Faculty Open Access Articles*. 1464. <https://digitalcommons.dartmouth.edu/facoa/1464>

This Article is brought to you for free and open access by Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Faculty Open Access Articles by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Authors

Natasha Grotz, Tama Fox, Erin Connolly, Walter Park, Mary Lou Guerinot, and David Eide

Identification of a family of zinc transporter genes from *Arabidopsis* that respond to zinc deficiency

NATASHA GROTZ*, TAMA FOX*, ERIN CONNOLLY*, WALTER PARK†, MARY LOU GUERINOT*‡, AND DAVID EIDES§

*Department of Biological Sciences, Dartmouth College, Hanover, NH 03755; †Department of Biochemistry and Molecular Biology, University of Minnesota-Duluth, Duluth, MN 55812; and §Nutritional Sciences Program, University of Missouri-Columbia, Columbia, MO 65211

Edited by Christopher R. Somerville, Carnegie Institution of Washington, Stanford, CA, and approved March 31, 1998 (received for review February 13, 1998)

ABSTRACT Millions of people worldwide suffer from nutritional imbalances of essential metals like zinc. These same metals, along with pollutants like cadmium and lead, contaminate soils at many sites around the world. In addition to posing a threat to human health, these metals can poison plants, livestock, and wildlife. Deciphering how metals are absorbed, transported, and incorporated as protein cofactors may help solve both of these problems. For example, edible plants could be engineered to serve as better dietary sources of metal nutrients, and other plant species could be tailored to remove metal ions from contaminated soils. We report here the cloning of the first zinc transporter genes from plants, the *ZIP1*, *ZIP2*, and *ZIP3* genes of *Arabidopsis thaliana*. Expression in yeast of these closely related genes confers zinc uptake activities. In the plant, *ZIP1* and *ZIP3* are expressed in roots in response to zinc deficiency, suggesting that they transport zinc from the soil into the plant. Although expression of *ZIP2* has not been detected, a fourth related *Arabidopsis* gene identified by genome sequencing, *ZIP4*, is induced in both shoots and roots of zinc-limited plants. Thus, *ZIP4* may transport zinc intracellularly or between plant tissues. These *ZIP* proteins define a family of metal ion transporters that are found in plants, protozoa, fungi, invertebrates, and vertebrates, making it now possible to address questions of metal ion accumulation and homeostasis in diverse organisms.

Zn is an essential catalytic component of over 300 enzymes, including alkaline phosphatase, alcohol dehydrogenase, Cu-Zn superoxide dismutase, and carbonic anhydrase. Zn also plays a critical structural role in many proteins. For example, several motifs found in transcriptional regulatory proteins are stabilized by Zn, including the Zn finger, Zn cluster, and RING finger domains. Proteins containing these domains are very common; the yeast genome sequencing project has determined that almost 2% of all yeast gene products contain these types of Zn binding domains (1, 2). Inside cells, Zn is neither oxidized nor reduced; thus, the essential roles of Zn in cells is based largely on its behavior as a divalent cation that has a strong tendency to form stable tetrahedral complexes (for a review see ref. 3).

Despite the importance of Zn as an essential micronutrient for plant growth, relatively few studies have examined the mechanisms and regulation of Zn absorption by roots. Zn is taken up from the soil solution as a divalent cation (4). Currently, there is little agreement as to whether uptake is via ion channels or via a divalent cation carrier protein and whether there is a link between uptake and metabolic energy transduction. Studies of Zn uptake in plants mainly have been focused on hyperaccumulators, i.e., plants that can grow in

soils containing high levels of Zn and accumulate high concentrations of Zn in their shoots. Certain populations of *Thlaspi caerulescens* can tolerate up to 40,000 μg of Zn g^{-1} tissue in their shoots whereas the normal Zn concentration for most plants is between 20 and 100 μg of Zn g^{-1} tissue (4). Radiotracer studies with *T. caerulescens* and a closely related nonhyperaccumulating species, *T. arvense*, suggested that Zn uptake is controlled by regulating the number of active transporters in the membrane (5). The V_{max} for the uptake of Zn was 4.5-fold greater for *T. caerulescens* than for the nonhyperaccumulator *T. arvense* whereas their K_m values were not significantly different. Once in the root, Zn is believed to be transported into the xylem, taken up by leaf cells, and then stored in the vacuoles of leaf cells thus preventing the buildup of toxic levels in the cytoplasm. These observations indicate that there are also Zn transporters that transport the metal between cells and into subcellular compartments within the plant. We describe here the successful isolation and characterization of Zn transporter genes from *Arabidopsis thaliana*. To our knowledge, these are the first zinc transporter genes to be identified in plants.

MATERIALS AND METHODS

Yeast Growth Conditions and Library Screening. Yeast cells were grown in 1% yeast extract, 2% peptone supplemented with 2% glucose (yeast extract/peptone/dextrose), and in synthetic-defined medium (6.7 g/liter yeast nitrogen base without amino acids) supplemented with 2% glucose, 0.1% casamino acids, 0.01% adenine, and 0.01% tryptophan. Media for plate cultures were prepared with 1.5% agar. Strain ZHY3 (*MAT α zrt1::LEU2 zrt2::HIS3 ade6 can1 his3 leu2 trp1 ura3*) (6) was transformed (7) with a plasmid library containing *A. thaliana* cDNAs inserted under the control of the yeast phosphoglycerate kinase promoter in pFL61 (8). The poly(A) RNA used to construct this library was obtained from whole young seedlings (stage two leaves). Transformants were inoculated directly onto 39 synthetic-defined agar plates at a density of $\approx 7,000$ transformants/plate. Because of the mutations in the *ZRT1* and *ZRT2* genes, synthetic-defined medium is zinc-limiting for ZHY3. Plasmid dependence of the zinc uptake effects was confirmed by selecting for transformants that had lost the plasmids by using 5-fluoroorotic acid (9).

DNA Manipulations and Sequence Analysis. DNA was prepared from each of the three yeast transformants, and the plasmids were transformed into *Escherichia coli* TOP10F' cells (Stratagene). Plasmid DNA was prepared and digested with *NotI*, and the insert fragments were cloned into *NotI*-digested Bluescript SK+ (Stratagene). Nucleotide sequences were ob-

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/957220-5\$2.00/0
PNAS is available online at <http://www.pnas.org>.

This paper was submitted directly (Track II) to the *Proceedings* office. Data deposition: The *ZIP1*, *ZIP2*, and *ZIP3* sequences reported in this paper have been deposited in the GenBank database (accessions nos. AF033535, AF033536, and AF033537, respectively).

‡To whom reprint requests should be addressed. e-mail: Mary.Lou.Guerinot@Dartmouth.edu.

tained on an Applied Biosystems model 373A Automated Sequencer (Dartmouth College Molecular Biology Core Facility, Hanover, NH) by using the Ready-reaction Dye-Terminator kit with AmpliTaq FS (Applied Biosystems). Computer database comparisons were performed by using BLAST (10); potential transmembrane domains were identified by using TOP-PREDII (11); and multiple sequence alignment was performed by using PILEUP and PRETTY (GCG). A cDNA version of the *ZIP4* coding sequence, constructed by using PCR, was cloned into pFL61 and used in yeast complementation and uptake assays.

Zinc Uptake Assays. Zinc uptake assays were performed as described for iron uptake (12) except that $^{65}\text{ZnCl}_2$ (Amersham) and low zinc medium-EDTA (13) were substituted for $^{59}\text{FeCl}_3$ and low iron medium-EDTA (14), respectively. Because of their different pH optima for uptake activity, the *ZIP1* and *ZIP3* activities were assayed at pH 4.7 and *ZIP2* activity was assayed at pH 6.0. Michaelis-Menten kinetic values were determined by using KINETASYST software (IntelliKinetics, State College, PA). Stock solutions of the chloride salts of cadmium, cobalt, copper, manganese, and nickel were prepared in distilled water at a concentration of 100 mM. A ZnCl_2 stock solution was prepared at 100 mM in 0.02 N HCl, and the FeCl_3 stock solution was prepared at 50 mM in 0.1 N HCl.

Plant Growth Conditions. Seeds of *A. thaliana* (Columbia *gl-1*) were surface-sterilized and plated on Gamborg's B5 medium (Sigma) (pH 5.8) containing 2% sucrose, 1 mM Mes, and 0.7% agar. Plates were kept at 4°C in the dark for 4 days before being incubated under constant light at 21°C for 10 days. Seedlings then were transferred to either zinc-sufficient or zinc-deficient hydroponic medium, formulated by using the GEOCHEM-PC program (15). This medium (pH 6.0) contains 2 mM $\text{Ca}(\text{NO}_3)_2$, 1 mM KNO_3 , 500 μM MgSO_4 , 80 μM KH_2PO_4 , 1 mM Mes, 10 μM H_3BO_3 , and 0.1 μM Na_2MoO_4 ; the micronutrients CuSO_4 , $\text{Fe}_2(\text{SO}_4)_3$, MnCl_2 , and NiCl_2 were added in the presence of 100 μM excess EDTA to give final concentrations of 10 μM CuEDTA , 50 μM FeEDTA , 0.6 μM MnEDTA , and 0.1 μM NiEDTA . For the zinc-sufficient medium, ZnEDTA was present at a final concentration of 10 μM . The seedlings then were grown hydroponically for an additional 10 days. The zinc status of the plants was confirmed by using inductively coupled argon plasma spectrometry.

Nucleic Acid Analysis. Poly(A) RNA was extracted from root and shoot fractions of plants grown either in zinc-sufficient or zinc-deficient media. Samples (1 μg) of RNA were denatured, electrophoresed on a 0.8% agarose/6.2% formaldehyde gel, and then transferred to a nylon membrane (Bio-trans, ICN). RNA was bound to the membrane by UV crosslinking (Stratalinker, Stratagene). The membrane was prehybridized, hybridized, and washed as described by Pilgrim and McClung (16) with the exception that washes were performed at 42°C. Membranes were stripped for reprobing with a boiling solution of 0.1% SDS. DNA fragments used as hybridization probes were radiolabeled by random priming (17).

RESULTS AND DISCUSSION

To identify potential zinc transporters from plants, an *A. thaliana* cDNA expression library was screened for clones that restored zinc-limited growth when expressed in a *Saccharomyces cerevisiae* *zrt1 zrt2* mutant. This mutant lacks both high and low affinity zinc uptake systems and is extremely sensitive to zinc limitation because of its reliance on less efficient uptake pathways (6). From the $\approx 270,000$ transformants obtained, three independent transformants were isolated in which the *zrt1 zrt2* phenotype was suppressed. Restriction mapping and Southern blot hybridization analysis of the plasmids obtained from these transformants demonstrated that cDNA clones of three different genes had been isolated. The entire cDNA

insert of each plasmid was sequenced. Comparison of these sequences with the current databases revealed that these cDNAs each encoded proteins similar to the products encoded by the *ZRT1* and *ZRT2* genes of *S. cerevisiae* and the *IRT1* gene of *A. thaliana* (18) (Fig. 1). For this reason, the genes were named *ZIP1*, *ZIP2*, and *ZIP3* (for ZRT, IRT-like Protein). *ZRT1* and *ZRT2* encode the high and low affinity zinc transporters in yeast, respectively (6, 13), and *IRT1* encodes a probable iron transporter (18). A fourth *ZIP* gene, *ZIP4*, was identified during sequencing of the *Arabidopsis* genome (GenBank accession no. U95973); given the similarity of its deduced protein product to those encoded by the other *ZIP* genes, we included *ZIP4* in subsequent analyses.

The ability of *ZIP1*, *ZIP2*, and *ZIP3* to suppress the growth defect of a *zrt1 zrt2* yeast mutant suggested that these genes may encode zinc transporters. To test this hypothesis, we examined ^{65}Zn accumulation in a *zrt1 zrt2* strain expressing each of these genes. At 0°C, no zinc accumulation was detected in *ZIP1*- or *ZIP3*-expressing strains (Fig. 2A and C) or in an untransformed control strain (data not shown). At 30°C, no zinc accumulation was detected in the untransformed control strain (ref. 6 and data not shown) whereas a high rate of accumulation was observed in the *ZIP1*- and *ZIP3*-expressing strains (≈ 270 and 710 fmol/min/ 10^6 cells, respectively). The rate of *ZIP1*- or *ZIP3*-dependent zinc accumulation was not altered greatly by variations in pH of the assay solution between 4.0 and 6.0 (data not shown). In contrast, *ZIP2*-dependent zinc accumulation was highly pH-dependent; no uptake activity was detectable in the *ZIP2*-expressing strain when assayed at pH values below 5.0 (data not shown), yet at pH 6.0, temperature-dependent zinc accumulation was observed (≈ 19 fmol/min/ 10^6 cells) (Fig. 2B).

A cDNA version of the *ZIP4* coding sequence, constructed by using PCR, was cloned into pFL61 and was used in yeast complementation and uptake assays. No uptake could be detected in yeast cells expressing the *ZIP4* gene nor could *ZIP4* complement the growth defect of the *zrt1 zrt2* mutant on zinc-limited medium. This lack of uptake may be due to subcellular rather than plasma membrane localization of *ZIP4* in yeast.

When assayed over a range of zinc concentrations, uptake activity dependent on expression of the *ZIP1*, *ZIP2*, and *ZIP3* genes was concentration-dependent and saturable (Fig. 2D-F). The Michaelis-Menten kinetic values derived from these data are described in Table 1. Of interest, the apparent K_m values of these transporters are similar to the levels of zinc commonly found in plant rhizospheres (19).

To assess whether metals in addition to zinc are substrates for *ZIP1*, *ZIP2*, or *ZIP3*, we tested other metal ions for their ability to inhibit zinc uptake mediated by these proteins. Zinc uptake by *ZIP1* was not inhibited by a 10-fold excess of Mn, Ni, Fe, or Co (Fig. 2G). Zinc was the most potent competitor, demonstrating that *ZIP1* prefers zinc as its substrate over these other metal ions. Cd and Cu also inhibited zinc uptake but to a lesser extent. Although the mechanism of these inhibitory effects is not yet known, these results suggest that Cd and Cu also may be substrates for *ZIP1*. Zinc uptake mediated by *ZIP3* was inhibited by Mn, Fe, Co, Cd, and Cu (Fig. 2I). Again, zinc had the greatest effect, indicating that *ZIP3* also prefers zinc. Cd and Cu inhibited zinc uptake mediated by *ZIP2* to as great a degree as zinc (Fig. 2H) suggesting that *ZIP2* also may have a high affinity for Cd and/or Cu. In summary, these three transporters show unique sensitivities to other metal ions that may reflect differences in their substrate specificities. We have found that expression of *ZIP1*, *ZIP2*, *ZIP3*, and *ZIP4* in a *fet3 fet4* mutant [DEY1453 (20)] failed to alter the iron-limited growth defect of this strain or alter its iron uptake activity. Thus, the *ZIP* proteins cannot transport iron.

Why does *Arabidopsis* have multiple zinc transporters? Zinc, like other metal ions, is transported from the soil into the root




Fig. 1. Alignment of the amino acid sequences of seven full length *Arabidopsis* and *Saccharomyces* ZIP proteins (IRT1, ZIP1, ZIP2, ZIP3, ZIP4, ZRT1, and ZRT2). The eight transmembrane domains predicted for each of these proteins are numbered I–VIII. Residues highlighted in blue are identical, and residues highlighted in green represent conservative substitutions. The histidines found in the variable region between transmembrane domains III and IV are highlighted in gold. Residues highlighted in pink are conserved histidines or acidic residues found within predicted transmembrane helices.

and then must cross both cellular and organellar membranes as it is distributed throughout the plant. Specific zinc transporters may play different roles in this process. Northern blot analysis demonstrated that *ZIP1* and *ZIP3* are zinc-responsive. These genes are expressed predominantly in the roots of zinc-deficient plants; little or no mRNA was detected in the

roots of zinc-sufficient plants or in the shoots of zinc-sufficient or zinc-deficient plants (Fig. 3). Despite repeated attempts, *ZIP2* mRNA could not be detected in plants grown under either condition. *ZIP4* also responds to zinc deficiency, but, unlike *ZIP1* and *ZIP3*, this gene is induced in both the shoots and roots of zinc-deficient plants. This result is especially




FIG. 2. Biochemical properties of zinc uptake in yeast expressing the *ZIP1*, *ZIP2*, or *ZIP3* genes. *zrt1 zrt2* (*ZHY3*) transformants expressing *ZIP1*, *ZIP2*, or *ZIP3* were grown to exponential phase ($\approx 5 \times 10^6$ cells/ml) in synthetic-defined medium and assayed for zinc uptake with ^{65}Zn . Because of their different pH optima for uptake activity, the *ZIP1* and *ZIP3* activities were assayed at pH 4.7, and *ZIP2* activity was assayed at pH 6.0. (A–C) Time and temperature dependence of zinc accumulation assayed with $10 \mu\text{M}$ (*ZIP1*, *ZIP3*) or $1 \mu\text{M}$ ^{65}Zn (*ZIP2*) at 0°C (open symbols) and at 30°C (closed symbols). (D–F) Concentration dependence of zinc uptake was determined by measuring zinc accumulation for 5 min over a range of substrate concentrations. Each point in A–F represents the mean of two experiments each performed in duplicate, and the error bars represent ± 1 SD. (G–I) Inhibition of ZIP-dependent uptake in yeast by other metals. Cells were assayed for zinc uptake rate with $10 \mu\text{M}$ (*ZIP1*, *ZIP3*) or $1 \mu\text{M}$ (*ZIP2*) ^{65}Zn in the absence (–) or presence of 10-fold excess of the chloride salts of the indicated metal ions. Each value represents the mean of four replicates. The asterisks indicate significant differences from control values ($P < 0.05$) as determined by one-way ANOVA followed by Scheffe's test.

interesting given the predicted targeting of *ZIP4* to chloroplasts (see below). The results of our Northern blot analyses are consistent with a role for *ZIP1* and *ZIP3* in the uptake of zinc from the rhizosphere and a role for *ZIP4* in the transport of zinc in plastids. It is not surprising that *ZIP* gene expression would be regulated; zinc is toxic when present in excess. We also examined *ZIP* gene expression in iron-sufficient and iron-deficient plants and saw no response of any of the *ZIP* genes to the same conditions of iron deficiency that induce *IRT1* expression (data not shown). Thus, *ZIP1*, *ZIP3*, and *ZIP4* respond specifically to a lack of zinc.

The predicted products of the *ZIP1*, *ZIP2*, *ZIP3*, and *ZIP4* genes are 355, 353, 339, and 374 aa in length, respectively. The *ZIP* proteins contain eight potential transmembrane domains, numbered I–VIII in Fig. 1. *ZIP1*, *ZIP2*, and *ZIP3* each have a potential signal sequence and are predicted to be plasma membrane proteins; *ZIP4* contains a potential chloroplast targeting sequence [PSORT, (21)]. To date, we have identified a total of 18 *ZIP* family members, including nine plant genes (including eight from *Arabidopsis* alone), two yeast genes, one gene in the protozoan *Trypanosoma brucei* variety *rhodesiense*, four genes in the nematode *C. elegans*, and two in humans. Family members range in length from 309 to 476 aa; this range

Table 1. Michaelis–Menten kinetic constants of zinc uptake in yeast

Gene	Apparent K_m , μM	V_{max} , fmol/min/ 10^6 cells
<i>ZIP1</i>	13 ± 2	693 ± 43
<i>ZIP2</i>	2 ± 0.3	107 ± 4
<i>ZIP3</i>	14 ± 2	3528 ± 207

is largely due to variation in the number of residues between transmembrane domains III and IV, a domain we have des-




FIG. 3. Regulation of *ZIP1*, *ZIP3*, and *ZIP4* mRNA levels by zinc availability. Gene probes were hybridized to a Northern blot containing $1 \mu\text{g}$ of poly(A) mRNA prepared from the roots or shoots of hydroponically grown plants. The same blot was used sequentially (after stripping) for hybridization to each of the gene probes.

ignated the "variable region." The amino acid sequences of all known ZIP family members were aligned, and a dendrogram describing their sequence similarities was generated (Fig. 4). The family can be divided into two subfamilies, one containing sequences from animals and the other containing the plant, fungal, and protozoan members. Within the plant/fungal/protozoan subfamily, the two zinc transporters from yeast form one group and all but one of the plant sequences form a second group.

A potential metal binding motif, containing multiple histidine residues, is found in the variable regions of almost all of the fully sequenced members of this family including IRT1, IRT2, ZIP1, ZIP4, ZRT1, and ZRT2. In contrast, ZIP2 and ZIP3 contain only a single histidine in this region. Studies of ZRT1 suggest that the histidine-rich motif is located on the intracellular face of the plasma membrane and also indicate that it is essential for transporter function (D.E. and M. Broderius, unpublished work). Three of the transmembrane domains (II, IV, and V) contain a histidine residue that is fully conserved among all family members. These histidines are predicted to lie on the polar face of amphipathic helices, suggesting a possible role for these residues in substrate transport through the membrane. Furthermore, conserved acidic amino acids present in the transmembrane domains of several of the ZIP proteins also may be important for substrate movement.

The ZIP genes of *Arabidopsis* represent four members of a rapidly growing family of eukaryotic proteins. To date, six of




FIG. 4. Dendrogram showing amino acid sequence relationships among the ZIP family members. GenBank accession numbers for sequences designated by name are as follows: IRT1, U27590; IRT2, T04324; ZIP4, U95973; IRT3, M35868; ZRT1, P32804; and ZRT2, X91258. The amino acid sequence of human 998569 is only available in the Entrez database. The *Oryza sativa* D49213, IRT2, IRT3, and *Homo sapiens* H20615 clones have not been sequenced completely.

the family members have been implicated in metal ion transport. These results demonstrate that the ZIP transporter family plays roles in metal ion metabolism in a diverse array of eukaryotic organisms. The ZIP family is structurally distinct from other metal ion transporters such as the CDF family (22), which includes the recently identified mammalian zinc effluxers P-type ATPases (23) that are involved in the transport of a variety of cations and the Nramp proteins recently implicated in divalent cation transport (24–26). The ZIP genes then offer a good starting point not only for understanding how metals cross membranes but also for engineering plant-based solutions to nutrient deficiencies and to environmental remediation.

We thank Ann Thering, Elizabeth Rogers, and Rob McClung for critical reading of the manuscript, Raad Gitan for analyzing ZIP4 in yeast, and Michael Rutzke for performing the inductively coupled argon plasma analyses. This work was supported by grants from the National Science Foundation to M.L.G. (IBN-9318093) and to D.E. (MCB-9405200).

1. Böhm, S., Frishman, D. & Mewes, H. W. (1997) *Nucleic Acids Res.* **25**, 2464–2469.
2. Schjerling, P. & Holmberg, S. (1996) *Nucleic Acids Res.* **24**, 4599–4607.
3. Berg, J. M. & Shi, Y. (1996) *Science* **271**, 1081–1085.
4. Marschner, H. (1995) *Mineral Nutrition of Higher Plants*, (Press Academic, Boston), 2nd Ed.
5. Lasat, M. M., Baker, A. J. M. & Kochian, L. V. (1996) *Plant Physiol.* **112**, 1715–1722.
6. Zhao, H. & Eide, D. (1996) *J. Biol. Chem.* **271**, 23203–23210.
7. Schiestl, R. H. & Gietz, R. D. (1989) *Curr. Genet.* **16**, 339–346.
8. Minet, M., Dufour, M. E. & Lacroute, F. (1992) *Plant J.* **2**, 417–422.
9. Boeke, J. D., Trueheart, J., Natsoulis, G. & Fink, G. R. (1987) *Methods Enzymol.* **154**, 164–175.
10. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* **215**, 403–410.
11. Claros, M. G. & Von Heijne, G. (1994) *Comput. Appl. Biosci.* **10**, 685–686.
12. Eide, D., Davis-Kaplan, S., Jordan, I., Sipe, D. & Kaplan, J. (1992) *J. Biol. Chem.* **267**, 20774–20781.
13. Zhao, H. & Eide, D. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 2454–2458.
14. Eide, D. & Guarente, L. (1992) *J. Gen. Microbiol.* **138**, 347–354.
15. Parker, D. R., Chaney, R. L. & Norvell, W. A. (1995) in *Chemical Equilibrium and Reaction Models*, eds. Loeppert, R. H., Schwab, A. P. & Goldberg, S. (Soil Society of America, Madison, WI), pp. 163–199.
16. Pilgrim, M. L. & McClung, C. R. (1993) *Plant Physiol.* **103**, 553–564.
17. Feinberg, A. P. & Vogelstein, B. (1984) *Anal. Biochem.* **137**, 266–267.
18. Eide, D., Broderius, M., Fett, J. & Guerinot, M. L. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 5624–5628.
19. Welch, R. M. (1995) *Crit. Rev. Plant Sci.* **14**, 49–82.
20. Dix, D. R., Bridgham, J. T., Broderius, M. A., Byersdorfer, C. A. & Eide, D. J. (1994) *J. Biol. Chem.* **269**, 26092–26099.
21. Nakai, K. & Kanehisa, M. (1992) *Genomics* **14**, 897–911.
22. Paulsen, L. T. & Saier, M. H. (1997) *J. Membrane Biol.* **156**, 99–103.
23. Axelsen, K. B. & Palmgren, M. G. (1998) *J. Mol. Evol.* **46**, 84–101.
24. Belouchi, A., Kwan, T. & Gros, P. (1997) *Plant Mol. Biol.* **33**, 1085–1092.
25. Fleming, M. D., Trenor, C. C., Su, M. A., Foerzler, D., Beier, D. R., Dietrich, W. F. & Andrews, N. C. (1997) *Nat. Genet.* **16**, 383–386.
26. Gunshin, H., Mackenzie, B., Berger, U. V., Gunshin, Y., Romero, M. F., Boron, W. F., Nussberger, S., Gollan, J. L. & Hediger, M. A. (1997) *Nature (London)* **388**, 482–488.