

Dartmouth College

Dartmouth Digital Commons

Open Dartmouth: Peer-reviewed articles by
Dartmouth faculty

Faculty Work

11-1996

Circadian Clock-Controlled Genes Isolated from *Neurospora Crassa* are Late Night- To Early Morning-Specific

Deborah Bell-Pedersen
Dartmouth College

Mari L. Shinohara
Dartmouth College

Jennifer J. Loros
Dartmouth College

Jay C. Dunlap
Dartmouth College

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Biochemistry Commons](#), [Genetics Commons](#), and the [Physiology Commons](#)

Dartmouth Digital Commons Citation

Bell-Pedersen, Deborah; Shinohara, Mari L.; Loros, Jennifer J.; and Dunlap, Jay C., "Circadian Clock-Controlled Genes Isolated from *Neurospora Crassa* are Late Night- To Early Morning-Specific" (1996). *Open Dartmouth: Peer-reviewed articles by Dartmouth faculty*. 1472.
<https://digitalcommons.dartmouth.edu/facoa/1472>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Peer-reviewed articles by Dartmouth faculty by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Circadian clock-controlled genes isolated from *Neurospora crassa* are late night- to early morning-specific

(development/*cgc*/*frq*/light)

DEBORAH BELL-PEDERSEN, MARI L. SHINOHARA, JENNIFER J. LOROS, AND JAY C. DUNLAP*

Department of Biochemistry, HB7200, Dartmouth Medical School, Hanover, NH 03755

Communicated by David D. Perkins, Stanford University, Stanford, CA, August 23, 1996 (received for review July 9, 1996)

ABSTRACT An endogenous circadian biological clock controls the temporal aspects of life in most organisms, including rhythmic control of genes involved in clock output pathways. In the fungus *Neurospora crassa*, one pathway known to be under control of the clock is asexual spore (conidia) development. To understand more fully the processes that are regulated by the *N. crassa* circadian clock, systematic screens were carried out for genes that oscillate at the transcriptional level. Time-of-day-specific cDNA libraries were generated and used in differential screens to identify six new clock-controlled genes (*cgc*s). Transcripts specific for each of the *cgc*s preferentially accumulate during the late night to early morning, although they vary with respect to steady-state mRNA levels and amplitude of the rhythm. Sequencing of the ends of the new *cgc* cDNAs revealed that *cgc-12* is identical to *N. crassa* *cmt* encoding copper metallothionein, providing the suggestion that not all clock-regulated genes in *N. crassa* are specifically involved in the development of conidia. This was supported by finding that half of the new *cgc*s, including *cmt* (*cgc-12*), are not transcriptionally induced by developmental or light signals. These data suggest a major role for the clock in the regulation of biological processes distinct from development.

Circadian rhythms, generated and controlled by an endogenous biological clock, are observed in a wide variety of organisms ranging from cyanobacteria to man (reviewed in ref. 1). These daily rhythms, which persist under constant environmental conditions, are observed at the biochemical, physiological, and behavioral levels. The essence of circadian rhythms involves the restriction of an activity to a particular time of day to coordinate biological processes with exogenous environmental cycles. This requires the regulation of genes involved in the output pathways by the circadian clock.

Lower eukaryotes provide powerful model organisms with which to study the molecular and biochemical bases of circadian rhythms. Included in this group are the fungi which harbor relatively simple circadian systems, where the clock can be studied in the absence of complex multicellular interactions. Owing to its long history of genetics and biochemistry, the circadian system of *Neurospora crassa* has been one of the most highly described (reviewed in refs. 2 and 3): genetic and molecular analyses have shown that *frq* encodes a central component of the *N. crassa* oscillator required for rhythmicity (reviewed in ref. 2). In *N. crassa* the clock controls the timing of asexual spore (conidia) development, where in a wild-type strain grown in constant darkness, conidiation is initiated once every 22 hr.

One useful approach for understanding clock control of cellular events involves the characterization of genes that are temporally regulated by the circadian clock. Systematic screens for clock-regulated genes have now been reported in *N. crassa* (4), *Synechococcus* (5), and *Drosophila* (6). In *N. crassa*, initial

screens for clock-regulated genes were carried out by using subtractive hybridization of morning versus evening RNAs (4). Two morning-specific clock-controlled genes, *cgc-1* [independently identified as a glucose-repressible gene (*grg-1*) (7) in *N. crassa* (4, 8)] and *cgc-2* [later shown to be allelic to *eas* (9, 10)] were identified, and transcripts were shown to cycle with periods reflecting the genotypes of the strains analyzed. Nuclear run-on experiments demonstrated that rhythmic expression of *cgc-1* and *eas* (*cgc-2*) results from clock control of transcription (11), implicating the involvement of cis-acting regulatory elements mediating temporal control. This was confirmed for the *eas* (*cgc-2*) gene through the localization of a critical positive-activating clock element that was shown to be both necessary and sufficient for clock-regulated expression (12). Additionally, the rhythm of conidiation is unaffected in both *cgc-1* and *eas* (*cgc-2*) null strains, demonstrating that both genes are true output genes (as opposed to clock genes) that do not feedback on the oscillator (8, 9).

The commitment of *N. crassa* to development, a commitment regulated in part by the circadian clock, signals a substantial restructuring of the organism accompanied by major changes in the repertoire of expressed genes (13–16). In fact, all of the circadian clock-controlled genes heretofore described in *N. crassa* are induced during development and by light, and are believed to be involved with the conidial developmental pathway. These include *eas* (*cgc-2*), which encodes a fungal hydrophobin, a component of the hydrophobic rodlet layer of conidia important for spore dissemination (9, 10, 17), and *cgc-1*, which is highly regulated during development and by light (8, 18). External factors such as carbon source limitation, desiccation, and light, known to induce development (13), also induce *al-3* (19) and the conidiation-associated genes *con-6* and *con-10*, all three of which seem to be circadianly regulated (14, 16, 19). Where data are available for all five of these clock-regulated genes, regulation by light, development, or by the clock appears to be independently conferred by separate elements acting at the level of each gene (10, 12, 14, 16–19) rather than through global coordinated regulation of a single factor by all three processes. This unanticipated one-to-one correspondence between circadian clock regulation and light/developmental regulation has prompted the notion that the sole output of the *N. crassa* clock might be associated with the developmental pathway leading to conidiation.

While the isolation of only two *cgc*s in the initial screen (4) implied that few genes are regulated by the clock in *N. crassa*, the subsequent discovery that these genes were highly expressed suggested that clock regulation might be more prevalent. In addition, since only two phases of the circadian cycle were examined, and clock control may be more extensive at

Abbreviations: CT, circadian time; *cgc*, clock-controlled gene; RFLP, restriction fragment length polymorphism; FGSC, Fungal Genetics Stock Center; CuMT, copper metallothionein.

*To whom reprint requests should be addressed. e-mail: jay.c.dunlap@dartmouth.edu.

other times of the day, many potential rhythmic genes were likely missed (4). For these reasons we initiated a new search for clock-controlled genes to gain a more complete understanding of the cellular processes that are temporally regulated and to see if processes other than development might be clock regulated.

In this study, time-of-day-specific cDNA libraries were generated and used in differential screens to identify six new *ccgs* in *N. crassa*. All of the *ccgs* display maximal transcript accumulation in the late night to early morning, and differ in overall levels, amplitude and absolute peak of accumulation. As compared with *ccg-1* and *eas(ccg-2)* which together represent $\approx 10\%$ of poly(A)⁺ species, the relative mRNA expression levels of the new *ccgs* are low. In addition, the *ccgs* are found not to be tightly linked, ruling out coordinate regulation of gene expression at specific chromosomal locations by the clock. Preliminary sequencing of the ends of the *ccgs* revealed that *ccg-12* is identical to the previously cloned copper metallothionein gene (*cmt*) of *N. crassa*, not known to be involved in development. Importantly, on further examination we found that half of the new *ccgs*, including *cmt (ccg-12)*, show no regulation in response to changes in light or factors that trigger development. This establishes that the *N. crassa* circadian clock may have substantial involvement in aspects of the life of the organism exclusive of the regulation of development.

MATERIALS AND METHODS

Strains and Growth Conditions. Strains of *N. crassa* used in this study include the *frq*⁺ strain 30-7 (*bd*; A), and the long period mutant 695-425 (*bd*; *frq*⁷; A); the band (*bd*) mutation enhances the circadian rhythm of conidiation (20) but does not affect the clock itself. Strains for restriction fragment length polymorphism (RFLP) mapping studies were obtained from the Fungal Genetics Stock Center (FGSC) (Kansas City). *Neurospora media* (Vogel's or Fries) and vegetative growth conditions were as described (21, 22). Bacterial strain XLI-Blue (Stratagene) was used for all phage and plasmid manipulations as described (23).

Culture Harvesting Conditions. For rhythmic RNA analyses, tissue was grown and synchronized by light to dark transfers as described (4). Light and developmental induction experiments were accomplished in strain 30-7 using published methods (12). Light treatments were carried out on cultures grown in the dark for 22 hr (CT12), a time of minimal expression for each of the *ccgs*.

cDNA Library Construction. Poly(A)⁺ RNA (2 μ g) isolated from *N. crassa* (24) was annealed to 0.6 μ g of oligo(dT)-*Xba*I primer-adaptor (Promega) by heating the reaction to 70°C for 3 min, and then cooling immediately on ice. First- and second-strand cDNA synthesis was carried out using the cDNA Synthesis System of BRL. Double-strand cDNA (ca. 250 ng) was ligated to 10 pM phosphorylated *Eco*RI adaptors (Promega), phenol extracted, and then digested with 10 units *Xba*I. The cDNA was size selected using a Sephacryl S-400 spin column (Promega) and ligated to λ Zap II (Stratagene). The phage were then packaged *in vitro* using Promega's Packagene extracts and plated on *Escherichia coli* XLI-Blue. Growth of the recombinant phage on media containing isopropyl β -D-thiogalactoside and 5-bromo-4-chloro-3-indolyl β -D-galactoside indicated that less than 1×10^4 plaque-forming units/ μ g of DNA were devoid of inserts.

Differential Screens and Plasmid Rescue. Differential screens of the time-of-day-specific cDNA libraries were accomplished by plating a total of 5×10^4 phage on XLI-Blue in a top agar overlay, and duplicate plaque lifts (23) were hybridized to 5×10^5 cpm of ³²P-labeled cDNA (specific activity, 5×10^7 cpm/ μ g) from the same and opposite time points. Following autoradiography, the signals from the duplicate lifts were compared by visual inspection.

In vivo excision of the pBluescript plasmid containing each *ccg* cDNA from the λ ZapII vector was accomplished by recircularization of the plasmid containing the cloned insert with M13 helper phage R408 as described in the Undigested Lambda ZapII Cloning Kit (Stratagene).

Nucleic Acid Isolation and Hybridization. RNA was isolated using either a large scale method (25) or a miniprep method (26) and Northern blots were processed as described (12). RNA loading was normalized to *rRNA* as indicated, which remains at constant levels under the growth conditions used (4). Densitometric data was acquired and analyzed as described (12).

RFLP Mapping and Sequencing. The multicent strains, comprising 38 individual progeny from a cross of Mauriceville and Oak-Ridge-derived strains were used in RFLP analysis. *N. crassa* genomic DNA was isolated from mycelia by the hexadecyltrimethylammonium bromide (CTAB) extraction method (27, 28). Restriction polymorphisms were detected by Southern blot analyses (23) of digested DNAs using the individual cDNAs as probes, with the exception of *ccg-8*. No polymorphisms were detected for the *ccg-8* gene using its corresponding cDNA probe, so a cosmid containing the *ccg-8* gene (X10:3G) was identified from an ordered cosmid library (Orbach and Sachs library, FGSC) and then used as a probe. The results were compared with published updates of RFLP maps in the *Fungal Genetics Newsletter* (29).

Dideoxy sequencing of the cDNA clones was initiated from the T7 and T3 sites in the *in vivo* excised pBluescript vector, using an Applied Biosystems Prism Dye-deoxy sequencing kit.

RESULTS

Generation of Time-of-Day-Specific cDNA Libraries. To identify genes under control of the circadian biological clock in *N. crassa*, time-of-day-specific cDNA libraries were constructed for use in differential screens. To enrich for the selection of the desired clock-controlled genes instead of nutritionally and developmentally regulated genes (that could fluctuate in the liquid culture conditions used to synchronize the clock), a clock mutant strain, *frq*⁷ was used. Unlike the 22-hr period in a *frq*⁺ strain, the *frq*⁷ strain has a period of 29 hr, and this mutation [a single base pair change in the FRQ protein (30)] is specific to the circadian clock (31, 32). In liquid cultures of *frq*⁺ versus *frq*⁷ strains, the time of harvest can be chosen such that the same time in real hours in constant darkness (DD) represents different circadian times (CTs) in the two strains. (CT represents the normalization of biological time to 24 circadian hours per cycle in strains or organisms with varying periods. By convention, CT0 represents dawn and CT12 represents dusk in a 12:12 light/dark cycle.) Thus, clock-regulated transcripts, responding to CT, will have cycled out of phase (as shown for the *ccg-1* gene at DD44 in Fig. 1A), whereas developmentally and nutritionally regulated transcripts, responding to sidereal time in culture, should remain at equivalent levels.

Poly(A)⁺ RNA was isolated from *frq*⁺ and *frq*⁷ cultures held for 37 hr [representing approximately CT6 (noon) in *frq*⁺ and approximately CT18 (midnight) in *frq*⁷] and 43 hr [representing approximately CT12 (dusk) in *frq*⁺ and approximately CT0 (dawn) in *frq*⁷] in constant darkness (Fig. 1A). To verify that the cultures were indeed rhythmic, RNA from each time point was hybridized to *eas(ccg-2)* and as expected, the message peaks in accumulation between CT18 and CT0 (Fig. 1B). Because the levels of *eas(ccg-2)* mRNA are not globally elevated in the *frq*⁷ strain (data not shown), the peaks at CT18 and CT0 are clearly due to clock control. The rhythmic mRNA from each of the four times was used to construct directional cDNA libraries in λ ZapII (see *Materials and Methods*). The total number of recombinant phage particles in each of the libraries exceeds 2.5×10^6 , indicating that each gene has a




FIG. 1. Isolation of RNA from *frq*⁺ and *frq*⁷ cultures representing four different times in the circadian day. (A) Densitometry of the clock-regulated *ccg-1* mRNA isolated from *N. crassa frq*⁺ (solid line) and *frq*⁷ (dotted line) cultures grown in continuous dark and harvested every 4 hr (4). The time in the dark (Hours DD) is indicated on the bottom of the plot, and the corresponding CT is shown on the top for both strains. Arrows point to the time of harvest (DD37 and DD43) used to isolate mRNA for the generation of the time-of-day-specific cDNA libraries. (B) Northern blot hybridization of total RNA (10 μg) extracted from *frq*⁺ and *frq*⁷ strains at DD37 and DD43 representing approximate CT18, CT0, CT6, and CT12 as shown in A. Hybridization is to the clock-controlled *eas(ccg-2)* gene.

greater than 99% chance of being represented in the library based on rough estimates of 2000 vegetative mRNA species in *N. crassa* (33).

As an internal control for the time-of-day-specific libraries, we analyzed the representation of both *ccg-1* and *eas(ccg-2)* by hybridizing a minimum of 1000 plaques from each of the libraries with probes specific for both genes (Table 1). Compatible with our rhythmic Northern blot analyses (this study and ref. 4), the *ccg-1* cDNA is most prevalent in the CT0 library, and the *eas(ccg-2)* cDNA is most prevalent in the CT18 and CT0 libraries. Additionally, we found *ccg-1* and *eas(ccg-2)* to be abundantly represented in the libraries, comprising 8.5% of the CT0 cDNAs and 2.8% of the CT18 cDNAs, respectively.

Identification of Clock-Controlled Genes by Differential Screens of the Time-of-Day-Specific Libraries. To identify clock-regulated genes, differential screens were performed with the cDNA libraries representing opposite phases of the circadian day. The screens were designed to identify messages specific to early morning (CT0 versus CT12), midday (CT6 versus CT18), early evening (CT12 versus CT0), and midnight (CT18 versus CT6).

Approximately 5 × 10⁴ phage from each library were hybridized to labeled cDNA from the same and opposite time points. Twenty eight plaques (out of ≈40,000 visualized signals) found to hybridize to cDNA of the same phase, but not the opposite phase, were chosen as candidate ccgs. Sixteen of these were shown by hybridization to be re-isolates of *ccg-1* and *eas(ccg-2)* (providing a convincing internal control) while the

Table 1. Representation of *ccg-1* and *eas(ccg-2)* in the time-of-day-specific libraries

Gene	CT0, %	CT6, %	CT12, %	CT18, %
<i>ccg-1</i>	8.5	0.9	0	0.3
<i>eas(ccg-2)</i>	2.1	0	0.3	2.8

Percentages are derived from the number of positive hybridization signals/total number of plaques probed.

other 12 clones (4-CT0, 1-CT6, 2-CT12, 5-CT18) represented candidate ccgs. When the candidate ccgs were hybridized to RNA from four different times of day (data not shown), hybridization signals were not detected for two clones, two did not display any apparent clock regulation, and one clone (*ccg-6*) was represented three times (as determined by preliminary sequencing of the ends of the remaining candidate cDNAs). In all, six new candidate ccgs emerged from this screen.

To confirm the circadian nature of the regulation of these genes, the six candidate ccgs were examined further by hybridization to RNA from both *frq*⁺ and *frq*⁷ cultures. Tissue was grown in constant darkness and harvested for RNA isolation every 4 hr over 2 consecutive days. A representative experiment is shown in Fig. 2 for the *ccg-4* gene. In the *frq*⁺ strain (Fig. 2A), the mRNA cycles and accumulates to peak levels at DD12 (CT1) in the first cycle and DD32–DD36 (CT0–4) in the second cycle with a wild-type period of about 22 hr. A similar rhythm is observed in the *frq*⁷ strain where the period is ≈29 hr (Fig. 2B). As predicted for a clock-regulated gene, the message cycles in both *frq*⁺ and *frq*⁷, with a period that is consistent with the genotype of the strain. This is particularly evident at DD36, where the mRNA levels are 180° out of phase in the two strains (Fig. 2C). Similar analyses were conducted on the other candidate genes, and the results of hybridizations to RNA from the *frq*⁷ strain are shown in Fig. 3. In these experiments, the same RNA (with the exception of *ccg-4*) is probed with each of the ccgs, and *eas(ccg-2)* is used as an internal control. Although each gene hybridizes to an mRNA displaying an oscillation with a period of ≈29 hr, distinct differences in individual patterns of regulation are evident.




FIG. 2. Criteria used to establish clock control of gene expression, as illustrated by the *ccg-4* gene. The steady-state levels of *ccg-4* mRNA were assayed by Northern blot analyses in *frq*⁺ (A) and *frq*⁷ (B) strains. Liquid cultures of mycelia were grown in constant darkness and harvested after the indicated times in the dark (Hours DD). The approximate CT at the time of harvests are shown below the autoradiograms. (C) Following autoradiography, *ccg-4* mRNA was quantitated by densitometry, and plotted as relative band intensity versus time in the dark for both *frq*⁺ (solid line) and *frq*⁷ (dotted line). Equal loading of the RNA was verified by inspection of ethidium bromide stained rRNA on the gel (data not shown)




FIG. 3. Rhythmic transcript accumulation of the *ccgs* in the *frq7* (29-hr period) strain. RNA was isolated from the *frq7* mutant strain at the indicated times (Hours DD) representing the approximate CTs shown at the top of the autoradiograms, and hybridized to RNA probes specific for each *ccg* (shown at left). Equivalent loading of RNA was verified by hybridization to *rRNA*. Exposure times of the autoradiograms are varied for visualization.

Transcript size, relative abundance, average peak of mRNA accumulation, and amplitude of the rhythm for each gene are summarized in Table 2. All of the genes show peak transcript accumulation around the late night (CT18-23) to early morning (CT0-5), ranging from CT18 (for *ccg-12*) to CT5 (for *ccg-4*); none peak in the late day (CT6-11) or early evening (CT12-17).

ccg-7, -8, and -12 Are Likely to be Involved in Clock-Output Pathways Distinct from Conidiation. To begin to characterize the *ccgs*, the ends of the cDNA clones were sequenced. Comparisons of partial cDNA sequence for the *ccg-12* gene (data not shown) in the data bases reveals that *ccg-12* is identical to the previously cloned *cmt* gene of *N. crassa* (34) encoding copper metallothionein (CuMT). CuMT (26 aa) is multifunctional, having a role in copper storage, detoxification, and metal transfer to copper containing proteins. RNA abundance levels of the *cmt* gene are regulated not only by the clock (these studies), but expression is also induced by copper

Table 2. Summary of clock-controlled genes

	Measured transcript size,* kb	Relative transcript abundance [†]	Average peak [‡]	Amplitude [§]
<i>ccg-1</i>	0.6	+++	CT3	×4
<i>eas(ccg-2)</i>	0.7	+++	CT22	×5
<i>ccg-4</i>	1.1	++	CT5	×11
<i>ccg-6</i>	0.9	+	CT19	×9
<i>ccg-7</i>	1.0	+	CT21	×4.5
<i>ccg-8</i>	2.2	–	CT20	×3.5
<i>ccg-9</i>	2.0	–	CT19	×4.5
<i>ccg-12</i>	0.5	+	CT18	×2

*Transcript sizes were determined by comparison to known molecular weight markers.

[†]Transcript abundance was roughly estimated from exposure times required to obtain a signal equivalent to the *eas(ccg-2)* transcript.

[‡]The average peak of transcript abundance was determined from densitometry of the RNA levels over the times analyzed from Fig. 3. The peak in message accumulation will vary slightly in different experiments.

[§]The amplitude of the rhythm was calculated from the data in Fig. 3 using densitometry, and is reported as half the maximal distance between the inflection levels.

ions, with maximal accumulation occurring 1 hr after the addition of copper ions to the growth media (34). Based on all previous studies of *N. crassa* clock-regulated genes, we anticipated that the new *ccgs* would be directly involved in conidiation; however, the finding that *ccg-12* encodes CuMT suggested otherwise, as no role for CuMT in conidiation has been reported. This prompted us to examine the transcripts from the *ccgs* for induction after conidiation is initiated (Fig. 4A).

The expression patterns of *ccg-4, -6, and -9* during conidial development are very similar to conidiation-specific genes (35), with each transcript accumulating to high levels about 4 hr after development is initiated by desiccation in constant light. In contrast, the abundance of the *cmt(ccg-12)* transcript remains unchanged, while both *ccg-7* and *ccg-8* show slightly higher levels of mRNA in undifferentiated vegetative hyphae (mycelia) (Fig. 4A; time 0).

Genes that are regulated during conidiation in *N. crassa* are in most, if not all, cases induced by light. Therefore, we also assayed the effects of light on transcript levels for the *ccgs* (Fig. 4B). Consistent with dual light and developmental control, the *ccg-4* and *ccg-6* mRNAs accumulate to higher levels after a light treatment at CT12 for 30 min (L30) as compared with the dark control (D30), and *ccg-9* mRNA accumulates to maximal levels after a 1-hr light treatment (L60). However, transcripts from *ccg-7, -8, or -12* do not display any light responsiveness at the times examined. In *N. crassa*, a light treatment at CT12 causes about a 2-hr phase delay (36); however, this delay cannot explain the observed light induction of the *ccgs* for two reasons. First, the response is rapid and not sustained over the short length of time examined. Second, the response to light varies for *ccgs* that peak at approximately the same time of day. In both experiments, the *ccg-4* transcript is found to be present at very low levels as compared with previous experiments (Figs. 2 and 3), suggesting that *ccg-4* expression may be sensitive to the differences in growth media or the developmental age in the different experimental protocols (see *Materials and Methods*). In addition, while only one transcript is observed for the *ccg-6* gene in the rhythmic Northern blot (Fig.




FIG. 4. Developmental and light regulation of the *ccgs*. (A) RNA isolated from tissue 0, 2, 4, and 8 hr after induction of conidiation was hybridized to each new *ccg* (shown on the left) and to *rRNA* to verify equivalent loading. (B) RNA was isolated from cultures grown for 22 hr in the dark (CT12) and harvested after an additional 30 min in the dark (D30), or transferred to the light for 30, 60, and 120 min (L30, L60, and L120, respectively). The RNA was hybridized to probes as indicated above.

of genes in the laboratory (unpublished data). This is similar to what has been reported in *Drosophila*, where roughly 8% of 261 head-specific cDNA clones examined hybridized to an oscillating mRNA with a greater than 2-fold amplitude in a 12 hr light/dark cycle (6). Together, these findings are in dramatic contrast to the apparent global control of gene expression by the clock observed in the cyanobacterium *Synechococcus* (5), suggesting that in eukaryotes comprehensive regulation of gene expression by the oscillator is not a general phenomenon.

Studies of circadian-regulated gene expression, both at the transcriptional and translational levels, in diverse organisms are beginning to identify important metabolic pathways regulated by the circadian clock. In many organisms, control of gene expression by the clock includes both day- and night-specific regulation (reviewed in refs. 41–44). However, of these genes, only the clock-regulated genes of *N. crassa* and *Drosophila* have been identified in an organism where a component of the circadian oscillator has been well characterized, and thus, can ultimately be placed in a genetic pathway from the clock to the rhythmic output genes.

N. crassa, an organism that has been well defined both biochemically and genetically, now provides a rich source of eukaryotic genes known to be under control of the circadian clock. The systematic isolation and characterization of clock-controlled genes provides the framework for future studies aimed at detailing the pathways and molecular components involved in the conveyance of temporal information from the clock to rhythmic cellular processes. Together, these analyses will provide an entrée to understanding the mechanisms by which circadian regulation of gene expression is accomplished. In the long term, this will enable comparisons among the components necessary to achieve clock-mediated rhythmic gene expression in *N. crassa* and other eukaryotes.

We wish to thank Nam-Hai Chua for suggesting the use of *frq*⁷. We also thank N. Garceau, S. Brody, and D. Ebbole for comments on the manuscript, and both past and present members of our laboratories for useful discussions. This work was supported by grants from the National Institute of Mental Health (MH01186 to J.C.D and MH44651 to J.C.D and J.J.L.), the Air Force Office of Scientific Research (F49620-94-1-0260 to J.J.L.), the National Science Foundation (MCB-9307299 to J.J.L.), the Norris Cotton Cancer Center core grant at Dartmouth Medical School, and by a National Institutes of Health postdoctoral fellowship (GM15185 to D.B-P.).

1. Edmunds, L. N. (1988) *Cellular and Molecular Bases of Biological Clocks* (Springer, New York).
2. Loros, J. J. (1995) *Semin. Neurosci.* **7**, 3–13.
3. Lakin-Thomas, P. L., Coté, G. G. & Brody, S. (1990) *Crit. Rev. Microbiol.* **17**, 365–416.
4. Loros, J. J., Denome, S. A. & Dunlap, J. C. (1989) *Science* **243**, 385–388.
5. Liu, Y., Tsinoremas, N. F., Johnson, C. H., Levedeva, N. V., Golden, S. S., Ishiura, M. & Kondo, T. (1995) *Genes Dev.* **9**, 1469–1478.
6. Van Gelder, R. N., Bae, H., Palazzolo, M. J. & Krasnow, M. A. (1995) *Curr. Biol.* **5**, 1424–1436.
7. McNally, M. T. & Free, S. J. (1988) *Curr. Genet.* **14**, 545–551.
8. Lindgren, K. M. (1994) Ph.D. thesis (Dartmouth Medical School, Hanover, NH).
9. Bell-Pedersen, D., Dunlap, J. C. & Loros, J. J. (1992) *Genes Dev.* **6**, 2382–2394.
10. Lauter, F.-R., Russo, V. E. & Yanofsky, C. (1992) *Genes Dev.* **6**, 2373–2381.
11. Loros, J. J. & Dunlap, J. C. (1991) *Mol. Cell. Biol.* **11**, 558–563.
12. Bell-Pedersen, D., Dunlap, J. C. & Loros, J. J. (1996) *Mol. Cell. Biol.* **16**, 513–521.
13. Springer, M. L. (1993) *BioEssays* **15**, 365–374.
14. Lauter, F.-R. & Yanofsky, C. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 8249–8253.
15. Schmidhauser, T. J., Lauter, F.-R., Schumacher, M., Zhou, W., Russo, V. E. A. & Yanofsky, C. (1994) *J. Biol. Chem.* **269**, 12060–12066.
16. Corrochano, L. M., Lauter F.-R., Ebbole, D. J. & Yanofsky, C. (1995) *Dev. Biol.* **176**, 190–200.
17. Arpaia, G., Loros, J. J., Dunlap, J. C., Morelli, G. & Macino, G. (1993) *Plant Physiol.* **102**, 1299–1305.
18. Arpaia, G., Loros, J. J., Dunlap, J. C., Morelli, G. & Macino, G. (1995) *Mol. Gen. Genet.* **247**, 157–163.
19. Arpaia, G., Carattoli, A. & Macino, G. (1995) *Dev. Biol.* **170**, 626–635.
20. Sargent, M. L., Briggs, W. R. & Woodward, D. O. (1966) *Plant Physiol.* **41**, 1343–1349.
21. Davis, R. L. & de Serres, F. J. (1970) *Methods Enzymol.* **17**, 79–143.
22. Metznerberg, R. L., Stevens, J. N., Selker, E. U. & Morzycka-Wroblewska, E. (1984) *Fungal Genet. Newsl.* **31**, 35–39.
23. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) in *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY), 2nd Ed.
24. Chirgwin, J. M., Przybyla, A. E., MacDonald, R. J. & Rutter, W. J. (1979) *Biochemistry* **18**, 5294–5299.
25. Reinert, W. R., Patel, V. B. & Giles, N. H. (1981) *Mol. Cell. Biol.* **1**, 829–835.
26. Yarden, O., Plamann, M., Ebbole, D. J. & Yanofsky, C. (1992) *EMBO J.* **11**, 2159–2166.
27. Zolan, M. E. & Pukkila, P. J. (1986) *Mol. Cell. Biol.* **6**, 195–200.
28. Taylor, J. W. & Natvig, D. O. (1987) in *Zoosporic Fungi in Teaching and Research*, eds. Fuller, M. S. & Jaworsky, A. (Southeastern, Athens, GA), pp. 252–258.
29. Metznerberg, R. L. & Grotelueschen, J. (1993) *Fungal Genet. Newsl.* **40**, 130–138.
30. Aronson, B. D., Johnson, K. A. & Dunlap, J. C. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 7683–7687.
31. Feldman, J. F. (1982) *Annu. Rev. Plant Physiol.* **33**, 583–608.
32. Feldman, J. F. & Dunlap, J. C. (1983) *Photochem. Photobiol. Rev.* **7**, 319–368.
33. Wong, L.-J. C. & Marzluf, G. A. (1980) *Biochim. Biophys. Acta* **607**, 122–135.
34. Munger, K., Germann, U. A. & Lerch, K. (1987) *Experientia Suppl. (Basel)* **52**, 393–399.
35. Berlin, V. & Yanofsky, C. (1985) *Mol. Cell. Biol.* **5**, 849–855.
36. Dharmananda, S. (1980) Ph.D. thesis (Univ. of California, Santa Cruz).
37. Perkins, D. D. (1992) *Fungal Genet. Newsl.* **39**, 61–70.
38. Aronson, B. D., Johnson, K. A., Loros, J. J. & Dunlap, J. C. (1994) *Science* **263**, 1578–1584.
39. Hardin, P., Hall, J. & Rosbash, M. (1993) in *Molecular Genetics of Biological Rhythms*, ed. Young, M. W. (Dekker, New York), pp. 155–169.
40. Sehgal, A. (1995) *Semin. Neurosci.* **7**, 27–35.
41. Kay, S. A. & Millar, A. J. (1993) in *Molecular Genetics of Biological Rhythms*, ed. Young, M. W. (Dekker, New York), pp. 73–89.
42. Cahill, G. M. & Besharse, J. C. (1995) *Prog. Retinal Res.* **14**, 267–291.
43. Sassone-Corsi, P. (1994) *Cell* **78**, 361–364.
44. Mittag, M., Lee, D.-H. & Hastings, J. W. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 5257–5261.