

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

5-11-2010

Noncommutative Topology and the World's Simplest Index Theorem

Erik van Erp
Dartmouth College

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Analysis Commons](#), and the [Geometry and Topology Commons](#)

Dartmouth Digital Commons Citation

van Erp, Erik, "Noncommutative Topology and the World's Simplest Index Theorem" (2010). *Dartmouth Scholarship*. 1494.

<https://digitalcommons.dartmouth.edu/facoa/1494>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Noncommutative topology and the world's simplest index theorem

Erik van Erp¹

Dartmouth College, Hanover, NH 03755

Edited by Richard V. Kadison, University of Pennsylvania, Philadelphia, PA, and approved March 6, 2010 (received for review January 29, 2010)

In this article we outline an approach to index theory on the basis of methods of noncommutative topology. We start with an explicit index theorem for second-order differential operators on 3-manifolds that are Fredholm but not elliptic. This low-brow index formula is expressed in terms of winding numbers. We then proceed to show how it is derived as a special case of an index theorem for hypoelliptic operators on contact manifolds. Finally, we discuss the noncommutative topology that is employed in the proof of this theorem. The article is intended to illustrate that noncommutative topology can be a powerful tool for proving results in classical analysis and geometry.

Introduction: Hypoelliptic Fredholm Index Theory

In Euclidean space, waves propagate freely in all directions. A standing wave with frequency ω is a solution of the eigenvalue problem of Laplace's equation,

$$-\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} - \frac{\partial^2 u}{\partial z^2} = \omega^2 u.$$

For any value of ω the solution space of this partial differential equation (PDE) is infinite-dimensional, consisting of superpositions of plane waves traveling in arbitrary directions. The degrees of freedom become severely restricted when we place the wave equation on a *closed manifold*. Suddenly, solutions exist only for a discrete set of eigenfrequencies ω , and even with the correct choice of ω the solution space is always finite-dimensional. The reason is intuitively obvious: Unlike the situation in R^3 , waves “wrap around” a closed manifold, and a standing wave can exist only if it wraps around in just the right way, always remaining exactly in phase with itself. This lucky coincidence is the exceptional case, and generically the local propagation law (the PDE) is in conflict with the obstruction presented by the global topology.

The index theorem of Atiyah and Singer is the ultimate expression of this connection between the local features of the PDE and the global topology of the manifold. The *index* of a partial differential operator P is not exactly the dimension of its solution space. One must correct this dimension by subtracting the number of linearly independent conditions that are required of the “source” v if the inhomogeneous equation $Pu = v$ is to have solutions. Thus, the analytic index of P is defined as

$$\text{Index } P = \dim \text{Kernel } P - \text{codim Range } P.$$

Atiyah and Singer found a cohomology class $[\sigma(P)]$ associated to the principal symbol (i.e., the highest-order part) of P and expressed the analytic index of P in terms of this cohomology class by means of a purely topological formula. This *topological index* can often be computed explicitly with the tools of algebraic topology, and finding a positive index may be the only way to prove that solutions to the PDE *exist*—which, as we have seen, is not the generic situation.

A limitation of the theorem of Atiyah and Singer is that it applies only to *elliptic* operators. Recall that whereas Laplace's equation is elliptic, the heat equation is parabolic, and the wave equation hyperbolic. What Laplace's equation and the heat equation (but not the wave equation) have in common is that all

solutions—even weak solutions in the sense of distribution theory—are smooth functions. Both equations are *hypoelliptic*. Hörmander pioneered a deep investigation of hypoelliptic equations. It is not true in general that every hypoelliptic operator on a closed manifold is a *Fredholm operator*, i.e., an operator with a well-defined and finite analytic index. But the methods typically used to prove hypoellipticity also imply Fredholmness. Given these facts, it seems natural to search for an index theorem along the lines of the Atiyah–Singer formula for important classes of hypoelliptic Fredholm operators that appear in the literature.

A positive indication that this is possible was obtained by Hörmander himself. In 1971, almost 10 years after Atiyah and Singer published their result, Hörmander showed that the formula of Atiyah and Singer applies to hypoelliptic operators of type (ρ, δ) with $0 \leq 1 - \rho \leq \delta < \rho \leq 1$ (1). In local coordinates, the full symbols of such operators are invertible outside a compact set, and these symbols can be glued together, by a partition of unity, to a well-defined class in $K^0(T^*M)$ to which the Atiyah–Singer formula applies.

Despite a growing literature devoted to the analysis of hypoelliptic operators, for over thirty years no progress was made in hypoelliptic index theory since the theorem of Hörmander. Only about a decade ago did Epstein and Melrose reopen the investigation (2, 3). Their goal was to derive an index formula for the index of certain hypoelliptic operators associated to *contact structures*. The context for their work was not Hörmander's theorem, but Boutet de Monvel's index theorem for *Toeplitz operators* on strictly pseudoconvex CR manifolds (4). Boutet de Monvel's theorem is essentially the Atiyah–Singer formula, except that the symbol of an elliptic operator is replaced by an expression derived from the symbolic calculus of the Toeplitz algebra.

Epstein and Melrose's goal was to extend Boutet de Monvel's formula to the *Heisenberg algebra* on a contact manifold. The Heisenberg algebra is a nonstandard pseudodifferential theory that extends the Toeplitz algebra and (like any pseudodifferential theory) includes also the algebra of differential operators. The work of Epstein and Melrose (unfortunately largely unpublished) is a veritable tour de force, and they made considerable progress toward a solution. Their final formula contains several factors that are not present in the formula of Atiyah and Singer and that are very hard to compute in practice. One of the key differences between the Atiyah–Singer and Boutet de Monvel formulas and the formula of Epstein–Melrose is that the latter is applied to a differential form (representing the operator) that is *not closed* and therefore not a proper cohomology class. We suspect that the failure to identify a cohomology class is the primary reason for the appearance of the various mysterious correction factors.

In the present article we discuss an approach to the hypoelliptic index that has its roots in *noncommutative topology*. As we will see, this modern perspective provides a satisfactory resolution of

Author contributions: E.v.E. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹E-mail: erik.van.erp@dartmouth.edu.

structure on M . By a foundational result in contact topology, every closed orientable 3-manifold admits a contact structure (Martinet's theorem). In fact, there exist contact structures ξ in every homotopy type of oriented 2-plane bundles in TM (Lutz's theorem). Thus, operators P of the type we are interested in exist on every orientable 3-manifold.

Orientability of M is a necessary condition, because the triple X, Y, Z provides a well-defined orientation for M (it has the same orientation as $Y, X, [Y, X] = -Z$). In what follows we make the unnecessary but simplifying assumption that ξ is (globally) oriented as well.

We now make an elementary but surprising observation.

Proposition 2. *The Z coefficient γ in the local presentations of the operator P is a well-defined global function $\gamma: M \rightarrow C$.*

Proof: Suppose that in some open set in M we have two alternative presentations of P as

$$\begin{aligned} P &= -X^2 - Y^2 + i\alpha X + i\beta Y + i\gamma Z + \delta \\ &= -A^2 - B^2 + i\alpha' A + i\beta' B + i\gamma' C + \delta, \end{aligned}$$

where $Z = [X, Y]$ and $C = [A, B]$ and where both X, Y and A, B are positively oriented. The second-order terms in the two representations can agree only if the pair A, B is obtained from X, Y by an orthogonal transformation

$$A = aX + bY, \quad B = cX + dY,$$

with

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SO(2).$$

Direct calculation shows that

$$\begin{aligned} A^2 + B^2 &= (a^2 + b^2)X^2 + (ab + cd)(XY + YX) + (c^2 + d^2)Y^2 \\ &\quad + (aX \cdot b + cX \cdot d)Y + (bY \cdot a + dY \cdot c)X \\ &= X^2 + Y^2 + \xi - \text{terms.} \end{aligned}$$

and similarly

$$\begin{aligned} C &= AB - BA = (ad - bc)(XY - YX) + (aX \cdot c - cX \cdot a \\ &\quad + bY \cdot c - dY \cdot a)X + (aX \cdot d - cX \cdot b + bY \cdot d - dY \cdot b)Y \\ &= Z + \xi - \text{terms.} \end{aligned}$$

Combining these results we see that $\gamma = \gamma'$.

To those accustomed to the Fredholm theory of *elliptic* operators, the following theorem will sound surprising.

Proposition 3. *The operator P is Fredholm if the range of the Z coefficient γ does not contain any odd integers.*

The calculus behind Proposition 3 further implies—not surprisingly once we accept the proposition—that the index of the operator P depends only on the contact structure ξ and the homotopy type of the function

$$\gamma: M \rightarrow C \setminus \{\dots, -5, -3, -1, 1, 3, 5, \dots\}.$$

These curious facts have been known since the late 1970s (see, for example, refs. 8 and 9). Beals and Greiner (6) devoted an entire monograph to the analysis of exactly these operators (in a slightly more general setting). But until the work of Epstein and Melrose

the extensive literature on hypoelliptic operators contains no mention of the following question: *Can we express the Fredholm index of P as a “topological index,” in terms of the homotopy type of the contact structure ξ and the map γ ?* To explain the gap of two or three decades between the analysis of these operators and the resolution of the associated index problem, recall that all *scalar elliptic* differential operators have zero index, as do *all elliptic* differential operators (acting on sections in a vector bundle) on odd-dimensional manifolds. What we have here is a scalar operator on an odd-dimensional manifold. If there was some obvious way to reduce the index problem for these hypoelliptic operators to an elliptic problem, one would expect to find a zero index in all cases. But now that the problem is solved we know that every 3-manifold (except for a homology sphere) admits hypoelliptic scalar differential operators of the type we are studying here that have nontrivial index.

The resolution of the problem can be stated very simply as follows. (A small detail needed to understand the theorem: The Euler class of an oriented contact 2-plane bundle is always *even*.)

Theorem 1. *Let L be an oriented link in M such that the 1-cycle $2[L] \in H_1(M, Z)$ represents the Poincaré dual of the Euler class $e(\xi) \in H^2(M, Z)$. For each odd integer k , the finite collection of loops $\gamma: L \rightarrow C \setminus \{k\}$ has a winding number*

$$\text{Ind}_{L,\gamma}(k) = \frac{1}{2\pi i} \int_L \frac{d\gamma}{\gamma - k} \in Z.$$

The Fredholm index of P is a Z -linear combination of these winding numbers,

$$\text{Index } P = \sum_{k \text{ odd}} k \cdot \text{Ind}_{L,\gamma}(k).$$

Observe that if all coefficients of P are real valued (as in Proposition 1), then γ is purely imaginary and the map $\gamma: M \rightarrow C \setminus \{\text{odds}\}$ is contractible. Theorem 1 implies that $\text{Index } P = 0$, which is not surprising and can probably be proven by a direct homotopy from P to a self-adjoint operator. However, Theorem 1 also implies that if the vector fields X and Y in the presentation

$$P = -X^2 - Y^2 + i\alpha X + i\beta Y + i\gamma Z + \delta$$

are *globally* defined, then also $\text{Index } P = 0$, regardless of the homotopy type of γ . I am not aware of an elementary proof of this corollary.

Changing Orders to Suit Our Needs

Before outlining the proof of Theorem 1 we must digress and explain the calculus behind Proposition 3. Why is it that Fredholmness of P depends on the *lower-order term* $i\gamma Z$? The answer, in a word, is that $i\gamma Z$ is included in the *highest-order part* of P , if only we redefine “order” in a suitable way. (References for the calculus described in this section are refs. 3, 6, and 10. However, the perspective taken here is slightly different and influenced by our work on index theory.)

The highest-order part of a differential operator like

$$P = \sum_{i,j=1}^3 a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^3 b_i \frac{\partial}{\partial x_i} + c$$

at a point $m \in M$,

$$P_m = \sum_{i,j=1}^3 a_{ij}(m) \frac{\partial^2}{\partial x_i \partial x_j},$$

is not well-defined as a differential operator on M . The algebra of differential operators is only filtered, not graded. Nevertheless, the highest-order part of P is well-defined and is usually taken to be a constant coefficient operator on the tangent fiber $T_m M \cong \mathbb{R}^3$, independent of coordinate choices.

All we need to do to include $i\gamma Z$ in the highest-order part of our hypoelliptic operators is to change the filtration on the algebra of differential operators. However, in so doing we also must reinvestigate what *kind* of object the highest-order part of an operator is.

The *Heisenberg calculus* is, in essence, nothing more than the working out of the consequences of an alternative filtration on the algebra of differential operators on M . The filtration is defined as follows: All vector fields in the direction of the contact field ξ (like X and Y) have ξ -order one, as usual, but any vector field that is not everywhere tangent to ξ (like Z) has ξ -order two. The highest-order part of P in the Heisenberg calculus will then clearly include the $i\gamma Z$ term.

But we must also investigate what “highest-order part” really means in this calculus. Abstractly, for any filtered algebra, the notion of highest-order part refers to an element in the associated graded algebra. So we need to find an analytic model for the associated graded algebra of the algebra of differential operators with the Heisenberg filtration.

Observe that in the associated graded algebra, smooth functions f commute with all vector fields A , because the commutator $[A, f] = A \cdot f$ is of order zero. It follows that elements in the graded algebra can be *localized* at points $m \in M$. The result of this localization is, formally,

$$P_m = -X(m)^2 - Y(m)^2 + i\gamma(m)Z(m).$$

As before, the highest-order part of P is not an operator on M , but it has a natural interpretation as a smooth family $P_m, m \in M$ of operators in the tangent fiber $T_m M$. The difference is that in the Heisenberg calculus the operator P_m is not a constant coefficient operator, but an operator that is *left invariant* for a nilpotent group structure on the tangent fiber. One identifies $T_m M$ (or, more precisely, $\xi_m \oplus T_m M / \xi_m$) with the *Heisenberg group*, by imposing the commutation relations

$$[X, Y] = Z, [X, Z] = [Y, Z] = 0.$$

Then the ξ -highest-order part of P is the smooth family $\{P_m, m \in M\}$ of left invariant homogeneous operators on the (graded) Heisenberg group

$$G_m = \xi_m \oplus T_m M / \xi_m \cong T_m M.$$

If one wants a more explicit formula for the model operator P_m , one could substitute

$$X(m) = \frac{\partial}{\partial x} - \frac{1}{2}y \frac{\partial}{\partial z}, \quad Y(m) = \frac{\partial}{\partial y} + \frac{1}{2}x \frac{\partial}{\partial z}, \quad Z(m) = \frac{\partial}{\partial z}.$$

Whereas this substitution is common in the analytic literature, it is not useful for our purposes. We prefer to retain the simpler algebraic expression of P_m in terms of X, Y , and Z .

We can now state the main results about hypoelliptic operators in the Heisenberg calculus.

Definition 4: A ξ -homogeneous invariant operator on a graded nilpotent group is a *Rockland operator* if $\pi(P_m)$ is invertible for all

irreducible unitary representations π of the group, except for the trivial representation.

Observe that for the Abelian group \mathbb{R}^n unitary representation theory amounts to Fourier theory, and a Rockland operator on \mathbb{R}^n is just a homogeneous elliptic constant coefficient operator. In this language, one could define ellipticity by saying that P is elliptic if each of the model operators P_m in the highest-order part of P is a Rockland operator on the Abelian group $T_m M$. This definition generalizes beautifully to the Heisenberg calculus.

Definition 5: A differential operator P is ξ -elliptic if all the model operators P_m in its ξ -highest order in the Heisenberg calculus are Rockland operators.

Proposition 6. Let (M, ξ) be a closed contact manifold, and P a scalar differential operator on M . If for all $m \in M$ the ξ -highest-order part P_m is a Rockland operator, then P is a hypoelliptic Fredholm operator.

Let us apply this to our example: When is $P_m = -X^2 - Y^2 + i\gamma(m)Z$ a Rockland operator? We need explicit formulas for the representations of the Heisenberg group. Because we are interested in the representation of the differential operator P_m , it is most convenient to express our formulas in terms of the Lie algebra. The Heisenberg group has two families of irreducible representations. First, there is an infinite family of scalar representations with two continuous parameters (x, y) , given by

$$\pi_{(x,y)}(X) = ix, \quad \pi_{(x,y)}(Y) = iy, \quad \pi_{(x,y)}(Z) = 0.$$

When we apply these to P_m , we find

$$\pi_{(x,y)}(P_m) = x^2 + y^2,$$

which is invertible unless $(x, y) = (0, 0)$ (the trivial representation). We see that invertibility of the nontrivial scalar representations of P_m amounts to “partial ellipticity” of P in the ξ -directions, and it is built into our definition of ξ from P .

The second family of irreducible representations of the Heisenberg group is labeled by a single parameter $t \in \mathbb{R} \setminus \{0\}$ ($t = 0$ corresponds to the entire collection of scalar representations). The representation space is the *Bargman–Fock* space. It is the completion of the space of holomorphic polynomials in one complex variable $z \in \mathbb{C}$. The monomials $z^q / q!$ form an orthonormal basis. The normalization is chosen such that the operators z and $\partial / \partial z$ are adjoints.

If $t > 0$, the representation π_t is defined by

$$\begin{aligned} \pi_t(X) &= i\sqrt{\frac{t}{2}} \left(z + \frac{\partial}{\partial z} \right), & \pi_t(Y) &= \sqrt{\frac{t}{2}} \left(z - \frac{\partial}{\partial z} \right), \\ \pi_t(Z) &= it, \end{aligned}$$

whereas π_{-t} is the conjugate representation of π_t . We find, for $t > 0$,

$$\begin{aligned} \pi_t(P_m) &= -\pi_t(X)^2 - \pi_t(Y)^2 + i\gamma(m)\pi_t(Z) \\ &= t \left(2z \frac{\partial}{\partial z} + 1 - \gamma(m) \right). \end{aligned}$$

As one easily verifies, the basis vectors $z^q / q!$ are eigenvectors for the operator $\pi_t(P_m)$ with eigenvalues $2q + 1 - \gamma(m)$. It follows that $\pi_t(P_m)$ is invertible for all $t > 0$ iff $\gamma(m)$ is not a positive odd integer.

Observe that, because of the *homogeneity* of P , we have

$$\pi_t(P_m) = t\pi_{+1}(P_m), \quad t > 0,$$

as can be seen explicitly in the formula above. Likewise,

$$\pi_{-t}(P_m) = t\pi_{-1}(P_m), \quad t > 0.$$

For this reason, it suffices to verify invertibility of $\pi_{+1}(P_m)$ and $\pi_{-1}(P_m)$. A convenient way of dealing with the conjugate representation π_{-1} is as follows. Consider the antiautomorphism of the Lie algebra of the Heisenberg group given by

$$X^{\text{op}} = X, \quad Y^{\text{op}} = Y, \quad Z^{\text{op}} = -Z.$$

This Lie algebra antiautomorphism induces an antiautomorphism of the algebra of invariant operators on G . Therefore invertibility of $\pi_{-1}(P_m)$ is equivalent to invertibility of $\pi_{+1}(P_m^{\text{op}})$. Thus, all we need to do is verify that $\pi_{+1}(P_m)$ and $\pi_{+1}(P_m^{\text{op}})$ are invertible. The advantage of this description is that we can work with the single representation π_{+1} , which will prove useful when we get to our index formula.

To finish the discussion, simply reversing the sign of γ in $\pi_{+1}(P_m)$ we obtain

$$\pi_{+1}(P_m^{\text{op}}) = 2z \frac{\partial}{\partial z} + 1 + \gamma(m).$$

This operator is invertible iff $\gamma(m)$ is not a *negative* odd integer. So, by Definition 5, P is ξ -elliptic precisely if $\gamma(m)$ is not an odd integer for any $m \in M$. Then Proposition 3, which seemed bizarre from the perspective of elliptic theory, appears as a special case of Proposition 6. Once appropriately generalized, what seemed strangely different is seen to be a perfect analogy.

The Atiyah–Singer Formula (sort of)

The previous discussion suggests that our hypoelliptic index problem should be rephrased, in the context of the Heisenberg calculus, as follows: *Find a topological index that expresses the Fredholm index of P in terms of its ξ -highest-order part*

$$\sigma_\xi(P) = \{P_m, m \in M\}.$$

This was, in essence, the problem that Epstein and Melrose set out to solve. Contrary to what is suggested by the complicated formula they found (announced in ref. 2), it turns out that the topological index for ξ -elliptic operators is no different from the topological index computed by Atiyah and Singer.

Theorem 2. (See ref. 11). *Let P be a ξ -elliptic differential operator on a closed contact manifold (M, ξ) . Then*

$$\text{Index } P = \int_M \text{Ch}(\sigma_\xi(P)) \wedge \text{Td}(M),$$

where

$$[\sigma_\xi(P)] = \left[\frac{\pi(P_m)}{\pi(P_m^{\text{op}})} \right] \in K^1(M).$$

This is the well-known cohomological formula of Atiyah and Singer, except that the principal symbol $\sigma(P)$ of an elliptic operator is replaced by the Heisenberg symbol $\sigma_\xi(P)$.

To explain our formula we need to clarify the meaning of the quotient

$$[\pi(P_m)\pi(P_m^{\text{op}})^{-1}] \in K^1(M),$$

which appears under the Chern character. Rather than explain this in general, we will illustrate its meaning by deriving Theorem 1 from Theorem 2 by explicitly computing the appropriate K -theory class.

So assume again that P is a ξ -elliptic scalar operator on a closed contact 3-manifold, locally presented as

$$P = -X^2 - Y^2 + i\alpha X + i\beta Y + i\gamma Z + \delta.$$

We have calculated $\pi_{+1}(P_m)$ and $\pi_{+1}(P_m^{\text{op}})$, and we see that the quotient of these two operators is a diagonal operator with eigenvalues

$$\frac{2q + 1 - \gamma(m)}{2q + 1 + \gamma(m)}.$$

These scalars act on the basis vector $z^q/q!$, which is canonically interpreted as a section in the q th tensor power $(\xi^{1,0})^{\otimes q}$. Here $\xi^{1,0}$ denotes the oriented 2-plane bundle ξ thought of as a complex line bundle.

Recall that, in general, a cocycle in $K^1(M)$ is represented by a pair $[E, \sigma]$ consisting of a vector bundle E on M and an automorphism σ of E . In the case of our ξ -elliptic operator P , we find

$$\left[\frac{\pi(P_m)}{\pi(P_m^{\text{op}})} \right] = \sum_{q=0}^{\infty} \left[(\xi^{1,0})^{\otimes q}, \frac{2q + 1 - \gamma}{2q + 1 + \gamma} \right] \in K^1(M).$$

The left-hand side appears in Theorem 2, and the right-hand side explains its meaning in our example. Observe that the right-hand side only *appears* to be an infinite sum, because for sufficiently large q the automorphisms are close to (and therefore homotopic to) the trivial automorphism.

It is this computation of a K -theory class associated to P that is unique to the Heisenberg calculus for contact manifolds. Once we have the right K -theory class, we can apply the formula of Atiyah and Singer. By using the fact that the Chern character is a ring homomorphism (from K theory to deRham cohomology), we find

$$\text{Ch}(\sigma_\xi(P)) = \sum_{q=0}^{\infty} \text{Ch}(\xi^{1,0})^q \wedge \text{Ch}\left(\frac{2q + 1 - \gamma}{2q + 1 + \gamma}\right).$$

Because $\xi^{1,0}$ is a complex line bundle on a 3-manifold, we have

$$\text{Ch}(\xi^{1,0})^q = (1 + c_1(\xi^{1,0}))^q = 1 + q \cdot e(\xi),$$

where $e(\xi)$ denotes the Euler class of the oriented real 2-plane field ξ . The (odd) Chern character of an invertible function $u: M \rightarrow \mathbb{C}$ is given by

$$\text{Ch}(u) = \frac{1}{2\pi i} d \log u.$$

Because $\dim M = 3$, we have $\text{Td}(\xi^{1,0}) = 1 + e(\xi)/2$, and the Atiyah–Singer formula gives

$$\begin{aligned} \text{Index } P &= \int_M \text{Ch}(\sigma_\xi(P)) \wedge \text{Td}(M) \\ &= \sum_{q=0}^{\infty} \int_M \frac{2q+1}{2} e(\xi) \wedge d \log(2q+1-\gamma) \\ &\quad - \frac{2q+1}{2} e(\xi) \wedge d \log(2q+1+\gamma), \end{aligned}$$

which is equivalent to Theorem 1.[†]

A Factory for Hypoelliptic Index Theorems

We have explained in what sense Theorem 1 is an application of the topological index of Atiyah and Singer. From the point of view of classical analysis or topology it is very surprising that the Atiyah–Singer formula can be modified in such a straightforward manner. Our next objective is to explain why this is so. As before, what appears strange from one point of view is clarified when seen from a higher level of abstraction. In the present case, we must understand the problem in the context of *noncommutative topology*—specifically, the theory of C^* algebras and analytic K theory.

The idea of analytic K theory is rooted in Atiyah’s representation of $K^0(X)$ for compact X as the set of homotopy classes of maps from X into \mathcal{F} ,

$$K^0(X) = [X, \mathcal{F}],$$

where \mathcal{F} is the space of all Fredholm operators on a given (infinite-dimensional, separable) Hilbert space. Concretely, if $\{P_x, x \in X\}$ is a continuous family of Fredholm operators P_x over X , then the family of kernels E_x of P_x defines a vector bundle E over X (possibly of nonconstant fiber dimension), and so does the family F_x of cokernels. The “index” of this family can then be thought of as the formal difference

$$[E] - [F] \in K^0(X).$$

This “family index” is Atiyah’s isomorphism.

The idea that K -theory elements can be represented by families of Fredholm operators has been generalized and developed into analytic K theory of C^* algebras. The limited space of this article does not allow us to explain these ideas here in any detail. However, the example of Atiyah’s theorem may help the reader accept the following facts without much explanation. We have seen that the highest-order part of an elliptic operator P on M is a family $\{P_m, m \in M\}$ of elliptic operators. These operators are not Fredholm, because they act on functions $C_c^\infty(T_m M)$ on the Euclidean space $T_m M$, which is not compact. But they define a “Fredholm operator” in a generalized sense. First, one can think of the family as a *single* differential operator on $C_c^\infty(TM)$ that differentiates only in the direction of the fibers. The *translation invariance* of the operators P_m can be expressed algebraically by treating $C_c^\infty(TM)$ as a *convolution algebra*, where functions on TM are multiplied by taking their convolution product in each fiber. Then the family $\{P_m, m \in M\}$ —taken as a single operator on $C_c^\infty(TM)$ —commutes with the right module action of $C_c^\infty(TM)$ on itself. Ellipticity of the individual operators P_m then implies that this $C_c^\infty(TM)$ -linear operator satisfies the axioms of a suitably generalized notion of Fredholm operator. One completes $C_c^\infty(TM)$ in a suitable norm to obtain the C^* algebra $C^*(TM)$. The $C^*(TM)$ -linear Fredholm operator $\{P_m, m \in M\}$ then defines an (unbounded) element in the analytic K theory of $C^*(TM)$,

[†]In a personal communication, Charles Epstein pointed out that the index formula for the special case of second-order differential operators can also be derived by a deformation of the Heisenberg symbol of the differential operator to the symbol of a Toeplitz operator, to which Boutet de Monvel’s index theorem applies. It is not clear if or how this trick extends to the general case.

$$[\{P_m, m \in M\}] \in K_0(C^*(TM)).$$

Of course, the Fourier transform in the fibers $T_m M$ of the tangent space gives an algebra isomorphism

$$C^*(TM) \cong C_0(T^*M).$$

After Fourier transform the family of operators $\{P_m, m \in M\}$ is replaced by the function $\sigma(P)$ on T^*M that is usually referred to as the “principal symbol”, and the analytic K -theory element $[\{P_m, m \in M\}]$ corresponds exactly to the topological K -theory class $[\sigma(P)]$ defined by Atiyah and Singer,

$$K_0(C^*(TM)) \cong K^0(T^*M) : [\{P_m, m \in M\}] \mapsto [\sigma(P)].$$

What is the point of this translation? After all, the definition of the topological class $[\sigma(P)]$ seems much simpler than that of the equivalent analytic class $[\{P_m, m \in M\}]$. It surely is. The point of the analytic construction is that it applies much more generally. In particular, it applies without essential modification to the highest-order part $\{P_m, m \in M\}$ of a ξ -elliptic operator P on a contact manifold. This time, the operators P_m are translation invariant for a nilpotent group structure in the fibers of TM . Let us denote by $T_\xi M$ the tangent space understood as the smooth family of Heisenberg groups $T_m M \cong \xi_m \oplus R$ that underly the Heisenberg calculus. As before, we can form the convolution algebra $C_c^\infty(T_\xi M)$ and its completion $C^*(T_\xi M)$. These algebras are noncommutative, but we do not have to worry about that. The highest-order part $\{P_m, m \in M\}$ of any operator P in the Heisenberg calculus can be conceived as an operator on $C_c^\infty(T_\xi M)$ that is right- $C_c^\infty(T_\xi M)$ linear. If, in addition, every P_m is a Rockland operator (which makes P ξ -elliptic), then one can prove that this family is indeed a Fredholm operator in the generalized sense of analytic K theory. Therefore, as before,

$$[\{P_m, m \in M\}] \in K_0(C^*(T_\xi M)).$$

So, at the very least, we have a K -theory element, albeit an analytically defined K -theory element. The goal is, then, to compute the corresponding topological K cocycle. Analytic K theory is as much noncommutative *topology* as it is analysis. In the present case, a noncommutative version of the Thom isomorphism in K theory (the Connes–Thom isomorphism) implies that

$$K_0(C^*(T_\xi M)) \cong K^0(T^*M).$$

Therefore, at least theoretically, our analytically constructed K -theory class can be interpreted as a topological K -theory class,

$$[\sigma_\xi(P)] \in K^0(T^*M).$$

In fact, the construction generalizes further. Suppose that M is equipped with an *arbitrary* subbundle $H \subseteq TM$, of arbitrary codimension. The bundle H could be a foliation, or a contact structure, it could be related to a sub-Riemannian structure on M , or it could be a random bundle with no geometric significance whatsoever. We can study hypoelliptic operators that are elliptic in the directions encoded by H and of lower order in the transversal directions. Such operators will be the elliptic elements in a generalized Heisenberg calculus associated to H (where transversal vector fields are given order two). The highest-order part for H -elliptic operators in this calculus is a family of invariant Rockland operators P_m on nilpotent groups $G_m = H_m \oplus T_m M/H_m$, and this family represents an analytic K -theory class

$$[\{P_m, m \in M\}] \in K_0(C^*(T_H M)).$$

Once again this group is naturally isomorphic to $K^0(T^*M)$, and so we have

$$[\sigma_H(P)] \in K^0(T^*M).$$

Thus, by using the tools of noncommutative topology, one can *always* construct a symbol class in $K^0(T^*M)$ for this type of hypoelliptic operators. Then a very general index problem presents itself: *Find the topological index*

$$\text{Ind}_H: K^0(T^*M) \rightarrow Z$$

for H -elliptic operators. A priori, this topological index could depend on the geometric structure H . But it does not. A beautiful argument involving deformation theory of C^* algebras (this is where Lie groupoids play a key role) implies that, once we have identified the symbol of a hypoelliptic operator as a class in $K^0(T^*M)$, there is only one topological index.

Theorem 3. *Let M be a closed manifold equipped with a subbundle $H \subseteq TM$. Let P be an H -elliptic operator on M —elliptic in the directions H and of lower order transversally. Then the highest-order part $\sigma_H(P) = \{P_m, m \in M\}$ in the generalized Heisenberg calculus for (M, H) defines a K -theory class*

$$[\sigma_H(P)] \in K_0(C^*(T_H M)) \cong K^0(T^*M),$$

and the Fredholm index of P is computed by the topological index of Atiyah and Singer,

$$\text{Index } P = \int_{T^*M} \text{Ch}(\sigma_H(P)) \wedge \text{Td}(M).$$

This theorem generalizes Boutet de Monvel's theorem for Toeplitz operators, and it is also the source of our proof of Theorem 2. Nevertheless, Theorem 3, by itself, is too abstract to be useful. To turn it into a computable formula—say, something as concrete as Theorem 1—two obstacles need to be overcome.

The first obstacle is a purely analytic problem in the Heisenberg calculus. Given an explicit differential operator P , it is not too hard to verify whether its H symbol consists of Rockland operators P_m . But that is not sufficient in a general setting. One must still prove that if each P_m is Rockland, then P has an inverse in the generalized Heisenberg calculus. This implication is known to be true if the manifold (with its structure H) can be locally identified with the graded nilpotent model group—such as is the case for foliations (Frobenius's theorem) and contact structures (Darboux's theorem). But in general it can be quite tricky. How hard the problem is in general can be gleaned by skimming through the monograph of Beals and Greiner (6). The bulk of that publication is devoted to proving just this fact for second-order H -elliptic operators in the case that H is a subbundle of codimension one in TM . To my knowledge, it is not known whether this fact (if all P_m are Rockland, then P has an inverse in the calculus) is true in absolute generality.

The second obstacle is a problem in noncommutative topology. In order to compute the Chern character of $\sigma_H(P)$, we must have an explicit understanding of the Connes–Thom isomorphism

$$K_0(C^*(T_H M)) \cong K^0(T^*M).$$

Unfortunately, there is no general method for finding a topological representation of the analytically defined class. Interestingly, the map from analytic to topological K theory is, in essence, a generalized index—as in the example of Atiyah's analytic representation of elements in $K^0(X)$. Therefore, this computation of the symbol class in $K^0(T^*M)$ is itself an index

problem—not the index of P , but the index of its highest-order part (which, as we have indicated, is a generalized Fredholm operator). The tractability of this problem depends on the specifics of the geometric structure (M, H) , as we will see in our next and final section.

Offspring: Foliations, Contact Manifolds, and More

To derive Theorem 2 from Theorem 3 and thereby give a proof of the world's simplest index theorem (Theorem 1), we need to find an explicit formula for the isomorphism

$$K_0(C^*(T_\xi M)) \cong K^0(T^*M)$$

if ξ is a contact structure. The computation depends on a careful analysis of the structure of the C^* algebra $C^*(T_\xi M)$.

In the general case, if $H \subseteq TM$ is an arbitrary bundle, the structure of the C^* algebra $C^*(T_H M)$ is clarified by the theory of *type I* C^* algebras. A C^* algebra is of type I if it can be decomposed, by a series of successive short exact sequences, into commutative C^* algebras or algebras that are K -theoretically equivalent to commutative algebras. This reduction to commutative algebras is precisely what establishes the connection between noncommutative and classical topology and will allow us to use topological K theory to compute an analytic K cocycle. Thus, the structure theory of C^* algebras tells us something significant about the geometric structure (M, H) and about the difficulty of the related index problem. It is reasonable to expect that the more terms there are in the decomposition series of $C^*(T_H M)$ (it is always finite), the more intractable the index problem will be.

In the trivial case where $H = TM$ —the case of elliptic operators—the computation is fairly straightforward, because of the isomorphism $C^*(TM) = C_0(T^*M)$ established by Fourier theory. In general, $C^*(T_H M)$ is commutative if the groups G_m are Abelian, which is the case precisely if $[H, H] \subseteq H$, i.e., if (M, H) is a *foliated manifold*. For foliations, the computation of the class $\sigma_H(P) \in K^0(T^*M)$ for H -elliptic operators (these are *not* elliptic) is not much harder than it is for elliptic operators. We have carried out this computation and by this general methodology derive a hypoelliptic index theorem that is similar in spirit to the one proved by Hörmander for hypoelliptic operators of type (ρ, δ) , except that hypoelliptic operators (and their parametrices) in the Heisenberg calculus are of type $(\frac{1}{2}, \frac{1}{2})$, a case not covered by Hörmander's theorem (see ref. 5).

From the perspective of the structure theory of $C^*(T_H M)$, contact structures are the next best thing, after foliations. If $H = \xi$ is a contact structure, there is a single short exact sequence that decomposes $C^*(T_\xi M)$, as follows:

$$0 \rightarrow C_0(M \times R^\times) \otimes \mathcal{K} \rightarrow C^*(T_\xi M) \rightarrow C_0(\xi^*) \rightarrow 0.$$

As before, we assume here that ξ is cooriented; i.e., TM/ξ is a trivial line bundle. Then $R^\times = R \setminus \{0\}$ refers to the punctured fiber of the dual of the line bundle $TM/\xi \cong M \times R$. The elementary algebra \mathcal{K} is the (abstract) C^* algebra of compact operators on a separable Hilbert space.

To understand this sequence, it is helpful to focus on a single fiber of the bundle of algebras $C^*(T_\xi M)$. For every $m \in M$, the fiber is the group C^* algebra $G^*(G_m)$ [the completion of the convolution algebra $C_c^\infty(G_m)$ in a suitable norm] of the Heisenberg group $G_m = H_m \times R$. This group algebra decomposes as

$$0 \rightarrow C_0(R^* \setminus \{0\}) \otimes \mathcal{K} \rightarrow C^*(G_m) \xrightarrow{\phi} C_0(H_m^*) \rightarrow 0.$$

The quotient map ϕ is obtained by assembling all the scalar representations of the group G_m , parametrized by points in H_m^* . The remaining irreducible representations of G are labeled by the dual of the center $R \subset G_m$ —excluding zero, which corresponds to the scalar representations. The decomposition

