

Dartmouth College

## Dartmouth Digital Commons

---

Open Dartmouth: Peer-reviewed articles by  
Dartmouth faculty

Faculty Work

---

8-16-1993

### Pseudostable Bubbles

Marcelo Gleiser  
*Dartmouth College*

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Physics Commons](#)

---

#### Dartmouth Digital Commons Citation

Gleiser, Marcelo, "Pseudostable Bubbles" (1993). *Open Dartmouth: Peer-reviewed articles by Dartmouth faculty*. 1983.

<https://digitalcommons.dartmouth.edu/facoa/1983>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Peer-reviewed articles by Dartmouth faculty by an authorized administrator of Dartmouth Digital Commons. For more information, please contact [dartmouthdigitalcommons@groups.dartmouth.edu](mailto:dartmouthdigitalcommons@groups.dartmouth.edu).

**PSEUDO-STABLE BUBBLES\***

Marcelo Gleiser

*Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755***Abstract**

The evolution of spherically symmetric unstable scalar field configurations (“bubbles”) is examined for both symmetric (SDWP) and asymmetric (ADWP) double-well potentials. Bubbles with initial static energies  $E_0 \lesssim E_{\text{crit}}$ , where  $E_{\text{crit}}$  is some critical value, shrink in a time scale determined by their linear dimension, or “radius”. Bubbles with  $E_0 \gtrsim E_{\text{crit}}$  evolve into time-dependent, localized configurations which are *very* long-lived compared to characteristic time-scales in the models examined. The stability of these configurations is investigated and possible applications are briefly discussed.

PACS : 11.10.Lm, 98.80.Cq, 64.60.-i

\* electronic mail: gleiser@peterpan.dartmouth.edu

A remarkable consequence of nonlinear field theories is the existence of localized, non-singular solutions of the classical equations of motion which are non-dissipative. In general, these solutions can be time-dependent or static. As is well-known, the existence and simplicity of static solutions is severely constrained by dimensionality [1]. For a single self-interacting real scalar field  $\phi(\mathbf{x}, t)$  such solutions are only possible in (1+1)-dimensions. More realistic (3+1)-dimensional static solutions must invoke more than one field, as in the case of the 't Hooft-Polyakov monopole [2].

Given their relevance to the study of nonperturbative effects in field theories, static, non-dissipative solutions have been, for the last twenty years or so, the focus of most efforts in the study of nonlinear solutions in classical field theories. However, within the last decade, the possibility that spontaneous symmetry breaking occurred in the early Universe has called for a better understanding of time-dependent phenomena in the context of relativistic field theories. For example, the dynamics of cosmological phase transitions [3] naturally invokes out-of-equilibrium conditions, with fields interacting with themselves and with a hot plasma in the background of an expanding Universe [4].

In the present work, the possibility that *time-dependent*, localized, non-dissipative solutions exist in the context of simple (3+1)-dimensional scalar field theories is examined. In particular, the focus will be on models involving only a single real scalar field with self-interactions dictated by a double-well potential. Since it is known that for a symmetric double-well potential (SDWP) all field configurations are unstable, it is possible to obtain the lifetime of a given spherically symmetric field configuration by numerically evolving the equation of motion. By adopting this procedure, it has been shown in the mid-seventies that certain configurations evolved into a state which was considerably long-lived (even though the lifetimes measured then were not very accurate). These time-dependent solutions were called “pulsons” [5]. However, not much has been done in order to further explore the properties of pulsons. A few exceptions, which are mainly related

to the existence of these solutions for the Sine-Gordon potential, different symmetries, and somewhat contrived stability studies, are listed in Ref. 6. In fact, recent studies of bubble evolution in SDWPs overlooked the existence of pulsons [7]. Also, their existence has never been investigated for ADWPs.

Here, it will be argued that the existence of pulsons is a very general feature of models with both symmetric *and* asymmetric potentials, depending only on the initial amplitude and energy of the configuration. It will always be assumed that the initial configuration interpolates between the two minima of the potential (bubbles with amplitude below a certain value cannot evolve into pulsons, as will be clear later), and that it is smooth enough, say,  $\phi(r, t = 0) \sim \exp[-r^2/R_0^2]$ , or  $\sim \tanh(r - R_0)$ , with  $R_0$  the initial “radius”. (From now on these configurations will be called the gaussian and the tanh-bubbles.) It is remarkable that during the nonlinear evolution of these configurations a regime of dynamical stability may be achieved in which energy is practically conserved within a localized region, due to some as yet unclear conspiracy between the many degrees of freedom involved.

The action for a real scalar field is,  $S[\phi] = \int d^4x \left[ \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V_{S(A)}(\phi) \right]$ , where the subscripts S and A stand for the SDWP and ADWPs, with,  $V_S(\phi) = \frac{\lambda}{4} \left( \phi^2 - \frac{m^2}{\lambda} \right)^2$ , and  $V_A(\phi) = \frac{m^2}{2} \phi^2 - \frac{\alpha}{3} m \phi^3 + \frac{\lambda}{4} \phi^4$ . A field configuration  $\phi_0(\mathbf{x}, t)$  is a solution of the equation of motion (an overdot denotes partial time derivative),  $\ddot{\phi} - \nabla^2 \phi = -\frac{\partial V(\phi)}{\partial \phi}$ , and has an energy,

$$E[\phi_0] = \int d^3x \left[ \frac{1}{2} \dot{\phi}_0^2 + \frac{1}{2} (\nabla \phi_0)^2 + V(\phi_0) \right]. \quad (1)$$

For a sufficiently well-localized spherically symmetric configuration  $\phi_0(r, t)$  (the case of interest here) with linear “size”  $\sim R_0$ , the integral over all space can be restricted to a spherical volume containing the configuration,

$$E[\phi_0](t) = E_K(t) + E_S(t) + E_V(t), \quad (2)$$

where the kinetic, surface, and volume energies are defined respectively by ( $\epsilon \geq 2$ )

$$E_K(t) = 2\pi \int_0^{\epsilon R_0} dr r^2 \dot{\phi}_0^2, \quad E_S(t) = 2\pi \int_0^{\epsilon R_0} dr r^2 (\phi_0')^2, \quad E_V(t) = 4\pi \int_0^{\epsilon R_0} dr r^2 V(\phi_0). \quad (3)$$

Introducing the dimensionless variables  $\Phi(r, t) = \sqrt{\lambda}\phi(r, t)/m$ ,  $\rho = m r$ , and  $\tau = m t$ , the equation of motion becomes, for both SDWP (and ADWP),

$$\frac{\partial^2 \Phi}{\partial \tau^2} - \frac{\partial^2 \Phi}{\partial \rho^2} - \frac{2}{\rho} \frac{\partial \Phi}{\partial \rho} = \Phi - \Phi^3 \quad (-\Phi + \tilde{\alpha} \Phi^2 - \Phi^3), \quad (4)$$

where  $\tilde{\alpha} \equiv \alpha/\sqrt{\lambda}$ . Note that for the ADWP, a solution where  $\Phi = 0$  is the local minimum is possible only if  $\tilde{\alpha}^2 > 9/2$ . The equation above was solved numerically using a finite differencing method fourth order accurate in space and second order accurate in time. Since the problem is two-dimensional a very fine grid could be used. By taking the lattice spacing to be  $h = 10^{-2}$  and the time step to be  $\theta = 5 \times 10^{-3}$ , energy was conserved throughout the evolution to one part in  $10^5$ .

Let us concentrate on the SDWP for now. In this case all bubbles are unstable, since there is no gain in volume energy in going from one vacuum to another. Take the vacuum to be at  $\Phi = -1$  and consider configurations which interpolate between the two minima. Both thick and thin-wall bubbles will be considered. In Fig. 1 the energy within a spherical shell, Eq. 2, is shown for initial configurations,  $\Phi_0(\rho, 0) = -\tanh(\rho - \rho_0)$ , with  $\rho_0 = 3$  and  $\rho_0 = 10$ , and for  $\Phi_0(\rho, 0) = 2\exp[-\rho^2/\rho_0^2] - 1$ , with  $\rho_0 = 4$  and  $\rho_0 = 8$ . Note the existence of an extended period of stability, of duration  $\sim 10^3 - 10^4 m^{-1}$ , where practically no energy is radiated away. Thus, a necessary condition (which was verified numerically) for the existence of this pseudo-stable behavior is, from Eq. 2

$$\frac{dE_K}{dt} \simeq - \left( \frac{dE_S}{dt} + \frac{dE_V}{dt} \right). \quad (5)$$

An interesting point is that the value of the energy at the plateau is fairly independent of the initial configuration. This suggests the existence of an attractor-like configuration

in field space which is approached in the course of the bubble's evolution. An extensive (but not exhaustive) search indicates that only for configurations of initial energies  $E_0 \gtrsim 60m/\lambda \equiv E_{\text{crit}}$  this behavior occurs. This corresponds to a gaussian bubble of radius  $R_0 \sim 2.4/m$ . For smaller energies, the evolution is well-fitted by the relation  $E(t) = E_0 \exp[-t/\tau_L(R_0)]$ , where  $\tau_L(R_0)$  is the bubble's lifetime. For gaussian bubbles of radii  $mR_0 = 1, 2, m\tau_L(R_0) = 3.5, 12$ , respectively.

From Fig. 1 it is clear that the evolution of large enough bubbles can be divided into three stages. First the bubble sheds its initial energy by quickly shrinking into a thick-wall bubble of energy roughly  $E \sim 50m/\lambda$ . For thin wall bubbles, like the tanh-bubble with  $\rho_0 \gg 1$ , this shrinking is well described by the relativistic motion of the bubble wall. (See, e.g., Ref. 7.) The field then settles into the pulson configuration. At any given time, the configuration is well-approximated by a half-gaussian, but with softer asymptotic behavior,  $\Phi(\rho \rightarrow \infty, \tau) \sim \exp(-\rho)$ . The field is localized within a small volume with linear size  $\sim 3/m$ , while its amplitude is rapidly oscillating about a value between the two minima. (In this author's opinion, a better name for these configurations would be "oscillons".) Finally, during the last stage of the evolution, the amplitude of oscillations decreases and the pulson quickly radiates its remaining energy away.

In order to gain some insight into the properties of the pulson consider the behavior of radial perturbations about  $\Phi_0(\rho, \tau)$ , defined as  $\Phi(\rho, \tau) = \Phi_0(\rho, \tau) + \delta\Phi(\rho, \tau)$ . Since  $\Phi_0(\rho, \tau)$  satisfies the equation of motion, expanding  $\delta\Phi(\rho, \tau)$  in normal modes,  $\delta\Phi(\rho, \tau) = \text{Re} \sum_n \phi_n(\rho) \exp(i\omega_n \tau)$ , it is found that the amplitudes  $\phi_n(\rho)$  satisfy the radial Schrödinger equation, (only the  $\ell = 0$  mode will be considered here),

$$-\frac{d^2 \phi_n}{d\rho^2} - \frac{2}{\rho} \frac{d\phi_n}{d\rho} + V(\rho, \tau_0) \phi_n = \lambda_n \phi_n, \quad (6)$$

where  $V(\rho, \tau_0) \equiv 3\Phi_0^2(\rho, \tau_0) - 3$ , and  $\lambda_n \equiv \omega_n^2 - 2$ .  $\lambda_n$  is introduced to make sure that  $V(\rho \rightarrow \infty) \rightarrow 0$ . The stability of a given configuration at time  $\tau_0$ ,  $\Phi_0(\rho, \tau_0)$ , is determined by the lowest eigenvalue being positive,  $\omega_0^2 > 0$ , or  $\lambda_0 > -2$ .

Clearly the general problem is quite complicated, as the potential  $V(\rho, \tau)$  is time-dependent. A reasonable simplification is to solve this equation for a given time  $\tau_0$  for which the field configuration  $\Phi_0(\rho, \tau_0)$  is obtained by evolving the equation of motion. This implicitly assumes that  $\Phi_0(\rho, \tau)$  varies slower than  $\delta\Phi(\rho, \tau)$ . Otherwise, unstable modes would not have enough time to grow and destabilize the pulson. One can then find the eigenvalue for a succession of snapshots and thus investigate its time evolution. The Schrödinger equation was solved using the shooting method [8]. To make sure the method worked, the equation was solved for the Coulomb potential and for the “kink potential”  $V(\rho) = 3\cosh^{-2}(\rho/\sqrt{2})$ , since in both cases the eigenvalues are known analytically [9]. (For the kink case, one must recall that the first eigenvalue in 3d corresponds to the second in 1d,  $E_0(3d) = E_1(1d) = -1/2$ , due to the boundary conditions at the origin.)

The results are shown in Fig. 2 during part of the pulson stage. Clearly, the pulson is stable against small radial perturbations. (No instability was detected during the whole pulson stage using the above method. An argument to explain the pulson’s final disappearance is advanced shortly.)

Next, the existence of pseudo-stable behavior for ADWPs is investigated. The asymmetry is controlled by the dimensionless coupling  $\tilde{\alpha}$ . As two examples, consider the nearly degenerate case,  $\tilde{\alpha} = 2.16$ , and the non-degenerate case,  $\tilde{\alpha} = 2.23$ . In the context of first-order phase transitions (finite temperature), a state initially localized at  $\Phi = 0$  will decay by the nucleation of bubbles larger than the critical bubble, which is an extremum of the  $O(3)$ -invariant Euclidean action. The critical bubble is thus the energy barrier for vacuum decay. For a given value of  $\tilde{\alpha}$  it is easy to obtain the critical bubble and its energy numerically, by using the shooting method. (Now, the method finds the value of the field in the bubble’s core.) For the two values of the asymmetry above, the energies and radii of the critical bubble are, respectively,  $E(\tilde{\alpha} = 2.16) \simeq 6.12 \times 10^2 m/\lambda$ ,  $R(\tilde{\alpha} = 2.16) \simeq 19.6/m$ , and  $E(\tilde{\alpha} = 2.23) \simeq 1.17 \times 10^2 m/\lambda$ ,  $R(\tilde{\alpha} = 2.23) \simeq 7.9/m$ . The

interest here is in investigating the evolution of *sub-critical* bubbles, to see if they display the pseudo-stable behavior observed for the SDWP. The results are shown in Fig. 3 where the energy within a spherical shell surrounding the initial configurations is shown as a function of time. For each value of  $\tilde{\alpha}$ , two tanh-bubbles with initial radii  $\rho_0 = 3$  and  $\rho_0 = 4$  were examined. Again, the existence of very long-lived pulsions is observed, with  $E_{\text{crit}}$  and lifetimes depending on the asymmetry. In Fig. 4 the phase-space portrait of the pulsion's core ( $\rho = 0$ ), for  $\tilde{\alpha} = 2.16$  and  $\rho_0 = 4$  is displayed for  $\tau \geq 1000$ . Note the similarity with the motion of a “damped” (an)harmonic oscillator. During the pulsion stage, the motion is restricted to a band in phase-space. As the pulsion becomes unstable, it will spiral around the final stable point  $\Phi = 0$  and  $\dot{\Phi} = 0$ ; As energy is gradually radiated away, the maximum possible amplitude (that is for  $\dot{\Phi} \simeq 0$ ) is driven below a critical value for stability. Bubbles with initial amplitudes below this value, roughly about the maximum of  $V(\Phi)$ , *and* small kinetic energy will quickly shrink. Note that this is the value below which the potential is approximately parabolic, so that nonlinear effects become subdominant. (Of course, small amplitude but high velocity bubbles can still escape the attractive well centered at  $\Phi = 0$ , as during the pseudo-stable pulsion stage.)

The present work raises many questions for future investigation. Apart from a more detailed study of the pulsion's properties and stability, it is still not clear why the initial configurations evolve into the pulsion stage. The pulsion's behavior is suggestive of some sort of nonlinear resonance effect occurring between the different modes of the field. It should be possible to separate the field into short and long wavelengths, with a cutoff around  $m^{-1}$ . The short wavelength modes would act as a perturbation, which due to the nonlinear coupling might induce the observed behavior. Apart from their potential interest for, among other topics, nonlinear optics and long Josephson junctions [10], these solutions are also of interest in the context of cosmological phase transitions [3]. Consider two possible regimes of interest, defined by the ratio of time scales  $\mathcal{R} = \tau_\phi/\tau_U \sim$



$(\phi/\dot{\phi})/(m_{Pl}/T^2) \sim T/m_{Pl}$ , where  $T$  is the temperature and  $m_{Pl}$  is the Planck mass. For  $\mathcal{R} \sim 10^{-3}$ , corresponding to the GUT scale, the extended lifetime of one sub-critical bubble can be relevant to the dynamics of a first-order transition. The Universe cools off quite fast and if the sub-critical bubble lives long enough it can *become* a critical bubble. For  $\mathcal{R} \sim 10^{-16}$ , corresponding to the electroweak scale, many sub-critical bubbles can be present before a critical bubble nucleates [11]. Given their long lifetimes, they could serve as nucleation sites for critical bubbles, very much like impurities in ordinary phase transitions, speeding the process of vacuum decay considerably. Finally, in the context of late-time phase transitions, it is desirable to have bubbles live longer in order for matter accretion to be efficient [12]. These and related questions are currently under investigation.

#### ACKNOWLEDGMENTS

I am grateful to J.D. Harris for many enlightening comments and to R. Ramos for useful discussions. I also thank M. Briggs for his numerical skills during the early stages of this investigation. This work is partially supported by a National Science Foundation grant No. PHYS-9204726.

## REFERENCES

1. G. H. Derrick, *J. Math. Phys.* **5**, 1252 (1964).
2. R. Rajaraman, *Solitons and Instantons*, North-Holland (1982).
3. E. W. Kolb and M. S. Turner, *The Early Universe*, Addison-Wesley (1990).
4. A few examples are, D. Boyanovsky and H. J. de Vega, *Phys. Rev.* **D47**, 2343 (1993); J. M. Cornwall and R. Bruinsma, *Phys. Rev.* **D38**, 3144 (1988); E. Calzetta and B. L. Hu, *Phys. Rev.* **D37**, 2878 (1988).
5. I. L. Bogolubsky and V. G. Makhankov, *Pis'ma Zh. Eksp. Teor. Fiz.* **24**, 15 (1976) [*JETP Lett.*, **24**, 12 (1976)]; *ibid.* **25**, 120 (1977) [*ibid.* **25**, 107 (1977)]; V. G. Makhankov, *Phys. Rep.* **C 35**, 1 (1978).
6. I. L. Bogolubsky, *Phys. Lett.* **A61**, 205 (1977); J. Geicke, *Phys. Scripta* **29**, 431 (1984); B. A. Malomed and E. M. Maslov, *Phys. Lett.* **A160**, 233 (1991); A. M. Srivastava, *Phys. Rev.* **D46**, 1353 (1992).
7. L. M. Widrow, *Phys. Rev.* **D40**, 1002 (1989).
8. W. H. Press, B. P. Flannery, S. A. Teukolsky, and W. T. Vetterling, *Numerical Recipes: The Art of Scientific Computing*, Cambridge University Press (1986).
9. L. D. Landau and E. M. Lifshitz, *Quantum Mechanics; nonrelativistic theory*, Addison-Wesley (1965).
10. For a review see, *Solitons*, Eds. S. E. Trullinger, V. E. Zakharov, and V. L. Prokovsky, North-Holland (1986).
11. M. Gleiser and E. W. Kolb, *Phys. Rev. Lett.* **69**, 1304 (1992).
12. See, for example, Z. Lalak and B. A. Ovrut, University of Pennsylvania Report No. UPR-0504T; C. Hill, D. Schramm, and J. Fry, *Comm. on Nucl. and Part. Phys.* **19**, 25 (1989).

## List of Figures

Figure 1. The energy within a spherical shell surrounding several initial configurations as a function of time for the SDWP. Larger bubbles have shorter lifetimes.

Figure 2. Time evolution of lowest eigenvalue. For clarity, points with  $\omega_0^2 > 2$  were displayed as  $\omega_0^2 = 2$ .

Figure 3. The energy within a spherical shell surrounding several initial configurations as a function of time for the ADWP.

Figure 4. Phase-space evolution of pulson's core for  $\tilde{\alpha} = 2.16$ ,  $\rho_0 = 4$ , and  $\tau \geq 1000$ .