

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

8-1-1985

Metabolic Effect of 3,3',5'-Triiodothyronine in Cultured Growth Hormone-Producing Rat Pituitary Tumor Cells. Evidence for a Unique Mechanism of Thyroid Hormone Action.

D L. St Germain
Dartmouth College

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Medicine and Health Sciences Commons](#)

Dartmouth Digital Commons Citation

St Germain, D L., "Metabolic Effect of 3,3',5'-Triiodothyronine in Cultured Growth Hormone-Producing Rat Pituitary Tumor Cells. Evidence for a Unique Mechanism of Thyroid Hormone Action." (1985). *Dartmouth Scholarship*. 3745.

<https://digitalcommons.dartmouth.edu/facoa/3745>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Metabolic Effect of 3,3',5'-Triiodothyronine in Cultured Growth Hormone-producing Rat Pituitary Tumor Cells Evidence for a Unique Mechanism of Thyroid Hormone Action

Donald L. St. Germain

Departments of Medicine and Physiology, Dartmouth Medical School, Hanover, New Hampshire 03756

Abstract

Physiologic levels of 3,3',5'-triiodothyronine (rT_3) are generally believed to have minimal metabolic effects in the pituitary gland and other tissues. In the present studies, the regulatory role of rT_3 and other thyroid hormones on iodothyronine 5'-deiodinase (I5'D) activity was studied in a growth hormone-producing rat pituitary tumor cell line (GH₃ cells). I5'D activity was thiol-dependent and displayed nonlinear reaction kinetics suggesting the presence of two enzymatic processes, one having a low Michaelis constant (K_m for thyroxine [T_4] of 2 nM) and a second with a high K_m value (0.9 μ M). Growth of cells in hormone-depleted medium resulted in a two- to 3.5-fold increase in low K_m I5'D activity ($P < 0.001$). The addition of thyroid hormones to the culture medium resulted in a rapid, dose-dependent inhibition of low K_m I5'D activity with the following order of analogue potency: $rT_3 \geq T_4 > 3,5,3'$ -triiodothyronine (T_3). Using serum-free culture conditions, rT_3 was ~50 times more active than T_3 . These inhibitory effects were noted within 15 min of hormone addition and could not be attributed to substrate competition with T_4 . These findings suggest that the control of T_4 to T_3 conversion by thyroid hormones in the anterior pituitary gland is mediated by a unique cellular mechanism that is independent of the nuclear T_3 receptor; and under some circumstances, rT_3 may play a regulatory role in controlling this enzymatic process.

Introduction

The metabolic actions of thyroid hormones appear to be mediated primarily through the interaction of 3,5,3'-triiodothyronine (T_3)¹ with specific nuclear receptors (1). In the various growth hormone-producing rat pituitary tumor cell

lines (GH cells) for example, T_3 is the most potent thyroid hormone analogue in stimulating glucose consumption (2), growth hormone production (3), and amino acid transport (4), and these effects correlate well with the proportion of nuclear binding sites occupied by T_3 . Although 3,3',5'-triiodothyronine (rT_3) also has agonist activity in GH cells, 1,000-fold greater concentrations are required, presumably due to the nuclear receptor having a significantly lower affinity for rT_3 (5). In other systems, the metabolic effects of rT_3 also appear to be minimal (5).

In the present studies, we used the GH₃ cell line to investigate the regulatory effects of thyroid hormones on a low Michaelis constant (K_m) iodothyronine 5'-deiodinase (I5'D) process (type II). Our finding that rT_3 is a potent and rapid inhibitor of this enzymatic process suggests that thyroid hormone control of T_3 formation in anterior pituitary tissue is mediated through a unique cellular mechanism that is independent of the nuclear T_3 receptor.

Methods

GH₃ cells were obtained from the American Type Culture Collection (Rockville, MD) and were routinely grown in 75-cm² plastic flasks and 100-mm plastic culture dishes using Ham's F-10 culture medium supplemented with 2.5% fetal calf serum (FCS), 15% horse serum (HS), and gentamycin (50 μ g/ml) in a humidified atmosphere of 5% CO₂/95% air. Medium was routinely changed twice weekly. In some experiments cells were grown to near confluence in standard medium and then transferred and maintained for 4 d in serum-free Ham's F-10 or in Ham's F-10 supplemented with FCS (10%) that previously had been treated with either charcoal (6) or an ion exchange resin (7). According to the manufacturer's (HyClone Laboratories, Logan, UT) specifications, the T_4 and T_3 levels were 12.7 μ g/dl and 118 ng/dl, respectively, in the FCS and 2.9 μ g/dl and 47 ng/dl, respectively, in the HS. Treatment of FCS either with charcoal or with an ion exchange resin was demonstrated to remove >99% of the iodothyronines.

Experiments were performed with cells at the near confluent stage of growth and 48 h after the last medium change. Cells were harvested by scraping with a rubber policeman, washed twice with ice-cold buffer (0.25 M sucrose, 0.02 M Tris-HCl, pH 7.6, 1 mM MgCl₂, 2 mM CaCl₂, 0.1 mM dithiothreitol [DTT], 5% glycerol), then resuspended in the same buffer and sonicated using a sonic dismembrator (Artek Systems Corp., Farmingdale, NY) for 25 s at a setting of 45. The sonicate was centrifuged at 800 g for 10 min and the supernatant was used immediately for determination of I5'D activity as previously described (8, 9). T_3 production rates were quantified by radioimmunoassay (RIA) using stable thyroxine (T_4) as a substrate at concentrations of 0.002–3.0 μ M in the presence of added DTT. The T_4 content of the supernatant fraction, before the addition of stable T_4 or enrichment with DTT, was determined by RIA in ethanol extracts (8). Protein concentration was determined by the method of Lowry et al. (10).

Portions of this work were presented at the 60th meeting of the American Thyroid Association, New York, September 1984.

Address correspondence to Dr. St. Germain, Division of Endocrinology, Dartmouth Medical School, Hanover, NH 03756.

Received for publication 28 February 1985.

1. *Abbreviations used in this paper:* DTT, dithiothreitol; FCS, fetal calf serum; GH cells, growth hormone-producing cell lines; HS, horse serum; I5'D, iodothyronine 5'-deiodinase; K_m , Michaelis constant; rT_3 , 3,3',5'-triiodothyronine; T_3 , 3,5,3'-triiodothyronine; T_4 , thyroxine.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/85/08/0890/04 \$1.00

Volume 76, August 1985, 890–893

Several experiments were performed to characterize and define the optimal conditions for quantifying I5'D activity in GH₃ cells: (a) enzyme activity in supernatants was found to be a linear function of both protein concentration (up to 11 mg/ml) and incubation time (up to 45 min) and was dependent on the presence of sulfhydryl reducing agents (maximum activity at 20 mM DTT). (b) No T₃ formation was noted in supernatants heated to 80°C for 45 min before incubation with T₄ and DTT. (c) Incubation of supernatants with known concentrations of stable T₃ (at 37°C for 45 min in the presence of 20 mM DTT) failed to disclose any evidence of T₃ degradation. (d) Enzyme activity determined 1–4 d after a medium change varied by <15% and was the same during either the log or near confluent stages of growth.

The uptake of rT₃ by GH₃ cells was determined by incubating cells in medium containing a known specific activity of ¹²⁵I-rT₃ (New England Nuclear, Boston, MA). Cells were then harvested, washed, and sonicated as above, and the amount of radioactivity in the supernate fraction was determined. The rT₃ content was calculated after correcting for the proportion of counts due to ¹²⁵I as determined by electrophoresis (11). Stable rT₃ was obtained from Calbiochem-Behring Corp., La Jolla, CA.

Kinetic data were analyzed using Eadie-Hofstee plots, and nonlinear plots were resolved into two components as previously described (9). Statistical comparison was performed using the unpaired *t* test. All results are given as mean±SE.

Results

I5'D activity in GH₃ cell supernatants demonstrated nonlinear reaction kinetics suggesting the presence of two enzymatic processes having *K_m* values of 2.4±1.0 nM (*n* = 4 experiments) and 0.9±0.3 μM (Fig. 1). Corresponding values for the maximum velocity were 23±7 fmol T₃/min mg protein and 32±6 fmol T₃/min mg protein. The addition of 0.5 mM 6-*n*-propyl-2-thiouracil to cell sonicates (in the presence of 5 mM DTT) inhibited T₃ production by <10% at low substrate concentrations (10 nM T₄) but by 53% (*P* < 0.001) at T₄ concentrations of 3 μM (data not shown). In subsequent experiments I5'D activity was determined using supernatant substrate concentrations of 10 or 50 nM T₄ and 20 mM DTT; these conditions permit quantification of primarily the low *K_m* process.

The effect on I5'D activity of culturing cells in medium supplemented with different serum preparations is shown in Fig. 2. Cells grown in Ham's F-10 medium supplemented with 10% FCS manifested only 46% of the activity noted in cells cultured in standard medium. However, cells maintained in serum-free medium or in the presence of 10% resin-treated or




Figure 1. Kinetic analysis, utilizing an Eadie-Hofstee plot, of I5'D activity in GH₃ cells. Cells were cultured in standard medium containing 2.5% FCS and 15% HS. Enzyme activity was determined using T₄ concentrations of 0.002–1.0 μM and 20 mM DTT. T₄ concentrations on the abscissa are expressed in nanomoles per liter. Points represent the mean of duplicate values, which differed <15%. Prot, protein.




Figure 2. The effect of different medium conditions on I5'D activity in GH₃ cells. Cells were grown to near confluence in standard medium (FCS/HS) and then transferred and maintained for four additional days in the medium indicated. I5'D activity was determined using a T₄ concentration of 10 nM and 20 mM DTT. Values represent the mean±SE of triplicate determinations. *, *P* < 0.001 compared with cells maintained in untreated FCS. Prot, protein.

charcoal-treated FCS showed a two- to 3.5-fold increase in I5'D activity, suggesting that one or more components of serum exerted inhibitory effects on enzyme activity. The addition of T₄, T₃, or rT₃ to the culture medium resulted in significant inhibitory effects on I5'D activity. Dose response curves, as determined in serum-free medium, are shown in Fig. 3. In this experiment, rT₃ was somewhat more active than T₄ and 50 times more active than T₃ in inhibiting I5'D activity. Additional experiments (*n* = 5) using both serum-free conditions and medium supplemented with resin-treated FCS confirmed that the order of potency of these analogues was rT₃ ≥ T₄ > T₃. In medium supplemented with resin-treated FCS, T₃ concentrations of >10^{−9} M were required to inhibit I5'D activity. Thyronine, at concentrations up to 10^{−7} M, inhibited this process by <10%.

The potential of rT₃ acting as a competitive substrate with T₄ in the reaction mixture was assessed by determining the rate of T₃ formation in the presence of both added T₄ (50 nM) and rT₃. Concentrations of rT₃ of up to 10 nM had no effect on T₃ production rates. Quantitation of rT₃ cellular uptake revealed that 4 h after the addition of 10^{−8} M rT₃ to the culture medium, supernatant concentrations of rT₃ were ~5 nM in cells grown in serum-free medium and 0.4 nM in cells maintained in medium supplemented with resin-treated FCS. Thus, even at high medium rT₃ concentrations, the supernatant concentrations of rT₃ were insufficient to affect T₃ production rates when 50 nM T₄ was provided as substrate in the reaction mixture. Furthermore, the concentration of T₄ (as determined by RIA) in supernatants derived from cells grown in medium supplemented with resin-treated FCS and either 10^{−8} M or 10^{−7} M T₄ were 0.7 nM and 7 nM, respectively; concentrations that would not add significantly to the rates of T₃ formation in the reaction mixture.²

2. Based on the measured sonicate protein concentration and the known values of protein content and cellular volume of GH cells (12), the actual intracellular concentrations of rT₃ and T₄ can be calculated. For cells incubated for 4 h in medium containing 10% FCS, intracellular rT₃ and T₄ concentrations were equal to the total medium concentrations of these iodothyronines. Under serum-free medium conditions, the intracellular concentrations were ~30-fold greater than the medium concentrations.




Figure 3. Dose response curves of the inhibitory effects of thyroid hormones on I5'D activity in GH₃ cells. Cells were grown to near confluence in standard medium and then transferred and maintained for four additional days in serum-free Ham's F-10 medium. Iodothyronines were added to the culture medium 4 h before harvesting. I5'D activity was determined using a T₄ concentration of 50 nM and 20 mM DTT. Points represent the mean of triplicate determinations, which differed <5%.

The regulatory effect of rT₃ on I5'D activity was shown to be quite rapid (Fig. 4). Significant inhibition was noted within 15 min of adding rT₃ to the medium and was near maximal at 2 h. A similar time course of inhibition was shown for T₄.

Discussion

I5'D activity in GH₃ cells was shown in the present studies to be a thiol-dependent process with nonlinear reaction kinetics suggesting the presence of two enzymatic pathways having markedly different K_m values for T₄. A differential sensitivity of these two processes to the inhibitory effects of PTU was also noted. These findings are similar to those previously described in homogenates of rat anterior pituitary glands (9, 13) and cerebral cortex (14). Our findings differ, however, from a prior report using GH₃ cells where only a high K_m process (3.8 μ M) was identified (15). The failure to detect a low K_m process in this latter study may have been secondary to the low concentrations of DTT used (2 mM) and the relative insensitivity of the T₃ RIA (3.75 pg/tube lower limits of detection vs. 0.2 pg/tube in the present studies [8, 9]).

Recent studies have shown that low K_m I5'D activity is




Figure 4. Time course of the inhibitory effect of 10⁻⁸ M rT₃ on I5'D activity in GH₃ cells. Cells were grown to near confluence in standard medium and then transferred and maintained in medium containing 10% resin-treated FCS. The rT₃ was added to the culture medium before harvesting at the times indicated. I5'D activity was determined using a T₄ concentration of 50 nM and 20 mM DTT. Points represent the mean of triplicate determinations, which differed <5%.

responsible for the vast majority of "local" T₃ formation in the pituitary and depends on thyroid hormone status (8, 13). In hypothyroid animals, the activity of this process is markedly elevated, whereas thyroid hormone administration induces a rapid decrease in T₄ to T₃ conversion. Our finding that low K_m I5'D activity in GH₃ cells rapidly responds to medium thyroid hormone concentrations supports the use of this cell line as a model system.

The effects of thyroid hormones on I5'D activity in various tissues are complex (16). In liver and kidney homogenates, rT₃ inhibits T₃ formation by substrate competition with T₄ for a high K_m (type I) I5'D process (17). The suppressive effects of thyroid hormones on low K_m (type II) I5'D activity in GH₃ cells, however, cannot be attributed to this mechanism. Our demonstration that rT₃ is a potent inhibitor of this process contrasts sharply with the weak agonist activity of this analogue on other metabolic processes (2, 5) and strongly suggests that a unique regulatory mechanism, independent of the nuclear T₃ receptor, is responsible for this effect. Further evidence for this is the finding that relatively high concentrations of T₃ (10⁻⁹ M in medium containing serum), which have been shown by other investigators to stimulate significantly growth hormone production and maximally suppress thyrotropin-releasing hormone receptors in GH₃ cells (18), had no effect on I5'D activity. A recent report (19) demonstrating that the injection of T₄ or rT₃ in the rat is more effective than T₃ in inhibiting pituitary and cerebral cortex I5'D activity, suggests that the order of iodothyronine potency found in GH₃ cells is also applicable *in vivo*. The cellular processes responsible for these regulatory effects are uncertain. Leonard et al. (20), however, have recently suggested that the T₃-dependent inhibition of I5'D activity in rat pituitary and cerebral cortex is mediated by a posttranscriptional mechanism that increases the rate of degradation or inactivation of the enzyme.

To date, a 5-deiodinase process has not been described in pituitary tissue and none could be demonstrated in the present study in GH₃ cells when T₃ was used as a substrate. This finding, together with the near equal potency of T₄ with rT₃, suggests that the regulatory effects of T₄ on I5'D activity are direct and not mediated by conversion to rT₃.

An intriguing question raised by these studies is the possible role of rT₃ in the control of T₄ to T₃ conversion in the pituitary and brain. Although rT₃ appears to be a potent regulator of this metabolic process, its free concentration in normal human serum is ~40-fold lower than that of T₄ (21). Thus, T₄ rather than rT₃ is likely to exert the principle physiologic control over this process. This thesis is consistent with reports that the administration of rT₃ to normal human subjects has no effect on the serum levels of T₄, T₃, thyrotropin, or the thyrotropin response to thyrotropin-releasing hormone (22, 23). In nonthyroidal illness, however, marked alterations occur in thyroid hormone metabolism, which may result in decreased serum free T₄ levels and increases in serum free rT₃ levels of up to 15-fold (21). Under such circumstances, inhibitory effects of rT₃ on I5'D activity may become significant.

Acknowledgments

The author would like to thank Dr. P. Reed Larsen for the generous gift of the T₃ and T₄ antiserum, Dr. Valerie Anne Galton and Dr. Lee A. Witters for support and advice, and Ms. Roberta McLain for technical assistance.

This research was supported by the National Institutes of Health Biomedical Support grant 2-SO7 RR 05392-22 and a grant from the Hitchcock Foundation (Hanover, NH). The author is the recipient of a Clinical Investigator Award (AM-01389).

References

1. Oppenheimer, J. H. 1979. Thyroid hormone action at the cellular level. *Science (Wash. DC)*. 203:971-979.
2. Samuels, H. H., J. S. Tsai, and R. Cintron. 1973. Thyroid hormone action: a cell-culture system responsive to physiologic concentrations of thyroid hormones. *Science (Wash. DC)*. 181:1253-1256.
3. Samuels, H. H., F. Stanley, and J. Casanova. 1979. Relationship of receptor affinity to the modulation of thyroid hormone nuclear receptor levels and growth hormone synthesis by L-triiodothyronine and iodothyronine analogues in cultured GH₁ cells. *J. Clin. Invest.* 63: 1229-1240.
4. Surks, M. I., E. C. Fels, and C. R. DeFesi. 1984. Induction of amino acid transport by L-triiodothyronine in cultured growth hormone-producing rat pituitary tumor cells (GC cells). *J. Biol. Chem.* 259: 5726-5733.
5. Papavasiliou, S. S., J. A. Martial, K. R. Latham, and J. D. Baxter. 1977. Thyroid hormonelike actions of 3,3',5'-L-triiodothyronine and 3,3'-diiodothyronine. *J. Clin. Invest.* 60:1230-1239.
6. Larsen, P. R. 1972. Direct radioimmunoassay of triiodothyronine in human serum. *J. Clin. Invest.* 52:1939-1949.
7. Samuels, H. H., F. Stanley, and J. Casanova. 1979. Depletion of L-3,5,3'-triiodothyronine and L-thyroxine in euthyroid calf serum for use in cell culture studies of the action of thyroid hormone. *Endocrinology*. 105:80-85.
8. St. Germain, D. L., and V. A. Galton. 1985. Comparative study of pituitary-thyroid hormone economy in fasting and hypothyroid rats. *J. Clin. Invest.* 75:679-688.
9. St. Germain, D. L., R. A. Adler, and V. A. Galton. 1985. Thyroxine 5'-deiodinase activity in anterior pituitary glands transplanted under the renal capsule in the rat. *Endocrinology*. 117:55-63.
10. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.
11. Galton, V. A. 1980. Binding of thyroid hormones in serum and liver cytosol of *Rana catesbeiana* tadpoles. *Endocrinology*. 107: 61-69.
12. Tashjian, A. H., Jr. 1979. Clonal strains of hormone producing pituitary cells. *Methods Enzymol.* 58:527-535.
13. Visser, T. J., M. M. Kaplan, J. L. Leonard, and P. R. Larsen. 1983. Evidence for two pathways of iodothyronine 5'-diiodination in rat pituitary that differ in kinetics, propylthiouracil sensitivity, and response to hypothyroidism. *J. Clin. Invest.* 71:992-1002.
14. Visser, T. J., J. L. Leonard, M. M. Kaplan, and P. R. Larsen. 1982. Kinetic evidence suggesting two mechanisms for iodothyronine 5'-deiodination in rat cerebral cortex. *Proc. Natl. Acad. Sci. USA*. 79: 5080-5084.
15. Melmed, S., M. Nelson, N. Kaplowitz, T. Yamada, and J. Hershman. 1981. Glutathione-dependent thyroxine 5'-monodeiodination modulates growth hormone production by cultured nonthyrotropic rat pituitary cells. *Endocrinology*. 108:970-976.
16. Larsen, P. R., J. E. Silva, and M. M. Kaplan. 1981. Relationship between circulating and intracellular thyroid hormones: physiologic and clinical implications. *Endo. Reviews*. 2:87-102.
17. Chopra, I. J. 1977. A study of extrathyroidal conversion of thyroxine (T₄) to 3,3',5'-triiodothyronine (T₃) in vitro. *Endocrinology*. 101:453-463.
18. Hinkle, P. M., M. H. Perrone, and T. L. Greer. 1979. Thyroid hormone action in pituitary cells: differences in the regulation of thyrotropin-releasing hormone receptors and growth hormone synthesis. *J. Biol. Chem.* 254:3907-3911.
19. Silva, J. E., and J. L. Leonard. 1985. Regulation of rat cerebrocortical and adenohypophyseal type II 5'-deiodinase by thyroxine, triiodothyronine, and reverse triiodothyronine. *Endocrinology*. 116: 1627-1635.
20. Leonard, J. L., J. E. Silva, M. M. Kaplan, S. A. Mellen, T. J. Visser, and P. R. Larsen. 1984. Acute posttranscriptional regulation of cerebrocortical and pituitary iodothyronine 5'-deiodinases by thyroid hormone. *Endocrinology*. 114:998-1004.
21. Kaptein, E. H., W. J. Robinson, D. A. Grieb, and J. T. Nicoloff. 1982. Peripheral serum thyroxine, triiodothyronine, and reverse triiodothyronine kinetics in the low thyroxine state of nonthyroidal illness: a noncompartmental analysis. *J. Clin. Invest.* 69:526-535.
22. Nicod, P., A. Burger, G. Strauch, A. G. Vagenakis, and L. E. Braverman. 1976. The failure of physiologic doses of reverse T₃ to effect thyroid-pituitary function in man. *J. Clin. Endocrinol. Metab.* 43:478-481.
23. Shulkin, B. L., and R. D. Utiger. 1984. Reverse triiodothyronine does not alter pituitary-thyroid function in normal subjects. *J. Clin. Endocrinol. Metab.* 58:1184-1187.