

Dartmouth College

## Dartmouth Digital Commons

---

Dartmouth Scholarship

Faculty Work

---

5-1-2015

### Sex-Specific Parental Effects on Offspring Lipid Levels

Irene M. Predazzi  
*Vanderbilt University*

Rafal S. Sobota  
*Dartmouth College*

Serena Sanna  
*Istituto di Ricerca Genetica e Biomedica (IRGB)*

William S. Bush  
*Vanderbilt University*

Jacqueline Bartlett  
*Dartmouth College*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Medicine and Health Sciences Commons](#)

---

#### Dartmouth Digital Commons Citation

Predazzi, Irene M.; Sobota, Rafal S.; Sanna, Serena; Bush, William S.; Bartlett, Jacqueline; Lilley, Jessica S.; Linton, Macrae F.; Schlessinger, David; Cucca, Francesco; Fazio, Sergio; and Williams, Scott M., "Sex-Specific Parental Effects on Offspring Lipid Levels" (2015). *Dartmouth Scholarship*. 3762.  
<https://digitalcommons.dartmouth.edu/facoa/3762>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact [dartmouthdigitalcommons@groups.dartmouth.edu](mailto:dartmouthdigitalcommons@groups.dartmouth.edu).

---

## Authors

Irene M. Predazzi, Rafal S. Sobota, Serena Sanna, William S. Bush, Jacqueline Bartlett, Jessica S. Lilley, Macrae F. Linton, David Schlessinger, Francesco Cucca, Sergio Fazio, and Scott M. Williams

## Sex-Specific Parental Effects on Offspring Lipid Levels

Irene M. Predazzi, PhD;\* Rafal S. Sobota, PhD;\* Serena Sanna, MS;\* William S. Bush, PhD; Jacqueline Bartlett, MS; Jessica S. Lilley, MD; MacRae F. Linton, MD; David Schlessinger, PhD; Francesco Cucca, MD; Sergio Fazio, MD, PhD; Scott M. Williams, PhD

**Background**—Plasma lipid levels are highly heritable traits, but known genetic loci can only explain a small portion of their heritability.

**Methods and Results**—In this study, we analyzed the role of parental levels of total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and triglycerides (TGs) in explaining the values of the corresponding traits in adult offspring. We also evaluated the contribution of nongenetic factors that influence lipid traits (age, body mass index, smoking, medications, and menopause) alone and in combination with variability at the genetic loci known to associate with TC, LDL-C, HDL-C, and TG levels. We performed comparisons among different sex-specific regression models in 416 families from the Framingham Heart Study and 304 from the SardiNIA cohort. Models including parental lipid levels explain significantly more of the trait variation than models without these measures, explaining up to  $\approx 39\%$  of the total trait variation. Of this variation, the parent-of-origin effect explains as much as  $\approx 15\%$  and it does so in a sex-specific way. This observation is not owing to shared environment, given that spouse-pair correlations were negligible ( $<1.5\%$  explained variation in all cases) and is distinct from previous genetic and acquired factors that are known to influence serum lipid levels.

**Conclusions**—These findings support the concept that unknown genetic and epigenetic contributors are responsible for most of the heritable component of the plasma lipid phenotype, and that, at present, the clinical utility of knowing age-matched parental lipid levels in assessing risk of dyslipidemia supersedes individual locus effects. Our results support the clinical utility of knowing parental lipid levels in assessing future risk of dyslipidemia. (*J Am Heart Assoc.* 2015;4:e001951 doi: 10.1161/JAHA.115.001951)

**Key Words:** cholesterol • genetics • lipids • risk factors • sex

Lipid levels are highly heritable traits, with estimates of 46% to 77% for total cholesterol (TC), 22% to 48% for triglycerides (TGs), 34% to 72% for low-density lipoprotein

cholesterol (LDL-C), and 37% to 82% for high-density lipoprotein cholesterol (HDL-C).<sup>1–3</sup> However, despite the recent improvements in technology and several large-scale genome-wide association (GWA) studies on this topic, the majority of the genetic contribution to lipid trait variation is still unexplained.<sup>1,3–11</sup> For example, a meta-analysis of cohorts including the Framingham Heart Study (FHS) was only able to explain 10% to 12% of total heritability in lipid concentrations when combining up to 95 relevant loci.<sup>11</sup> Several explanations have been proposed for the missing heritability of traits, such as lipid levels, including gene-gene and gene-environment interactions, rare variants not detected in the large GWA studies, or epigenetic influences not assessed in traditional genetic studies.<sup>10,12,13</sup> Based on previous observations that maternal environment influences cardiovascular (CV) outcomes in adult offspring,<sup>14–20</sup> that genetic associations could be sex-specific,<sup>3</sup> and that parent-of-origin effects (POE) influence several traits in animal models,<sup>21–26</sup> we hypothesized that (1) parental lipid traits explain a significant amount of the offspring lipid variation that is not accounted for by known genetic variants and (2) the effects of parental lipid traits are sex-specific.<sup>3</sup>

From the Atherosclerosis Research Unit, Departments of Medicine and Pharmacology (I.M.P., J.S.L., M.F.L., S.F.) and Center for Human Genetics Research (R.S.S., W.S.B.), Vanderbilt University Medical Center, Nashville, TN; Knight Cardiovascular Institute, Center for Preventive Cardiology, Oregon Health and Science University, Portland, OR (I.M.P., S.F.); Department of Genetics, Geisel School of Medicine, Dartmouth College, Hanover, NH (R.S.S., J.B., S.M.W.); Istituto di Ricerca Genetica e Biomedica (IRGB), CNR, Monserato, Italy (S.S., F.C.); Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH (W.S.B.); Division of Endocrinology, Department of Pediatrics, University of Mississippi School of Medicine, Jackson, MS (J.S.L.); Laboratory of Genetics, NIA, Baltimore, MD (D.S.).

An accompanying Tables S1 through S16 is available at <http://jaha.ahajournals.org/content/4/7/e001951/suppl/DC1>

\*Dr Predazzi, Dr Sobota, and Dr Sanna contributed equally to the work.

**Correspondence to:** Sergio Fazio, MD, PhD, Knight Cardiovascular Institute, Center for Preventive Cardiology, Oregon Health and Science University, 3181 SW Sam Jackson Park Rd, Portland, OR 97239. E-mail: [fazio@ohsu.edu](mailto:fazio@ohsu.edu)

Received February 25, 2015; accepted May 1, 2015.

© 2015 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

To test our hypotheses, we assessed the parent-offspring relationship of lipids in trios from 2 large, well-characterized cohorts: the FHS Offspring cohort and the SardiNIA cohort. Both studies include subjects of European ancestry and contain data from multiple generations.

## Methods

### Study Participants

The FHS is a prospective cohort originally designed to assess the epidemiology of CV disease (CVD).<sup>27,28</sup> Data have been collected from 3 generations of participants since its inception in 1948. The original cohort involved 5209 participants, 5124 were enrolled in the second generation starting in 1971, and 4095 in the third generation starting in 2002.<sup>27,28</sup>

Our analysis included participants from the second and third generations of the FHS for whom serum TC, TG, LDL-C (calculated using the Friedewald equation:  $\text{LDL-C} = \text{TC} - \text{HDL} - (\text{TG}/5)$ ) and HDL-C were available for both generations.<sup>29</sup> To limit confounding of results by relatedness, we only considered the oldest offspring for each nuclear family, creating parent-offspring trios. The final study population consisted of 416 trios, with 228 females and 188 males in the offspring generation.

The SardiNIA study is a longitudinal study designed to assess the epidemiology and genetics of aging-associated conditions.<sup>30</sup> The study enrolled 6921 volunteers from a cluster of 4 towns on the east coast of Sardinia and represents a collection of large pedigrees, containing data from up to 5 generations. From each pedigree, we only considered parents and their oldest offspring from the 2 most recent generations.

The SardiNIA cohort had 304 families that were included in the analyses. Because this cohort included several families with only a single parent enrolled, analyses were performed on 277 mothers, 151 fathers, 168 daughters, and 136 sons. This study population included 124 complete parent-offspring trios.

For both cohorts, families in which any member had a TG level >400 mg/dL were excluded. Only individuals with at least 2 of the 4 lipid traits available were included in the analysis.

The study was approved by the institutional review boards at Vanderbilt University, Boston University, Dartmouth College, the National Institutes of Health, and the local ethical committee of Lanusei, Sardinia, in Italy. All of the included FHS and SardiNIA participants provided written informed consent, including consent to use of their DNA data in genetic analyses. For both cohorts, false paternity was assessed by genetics data management groups and the parental informa-

tion was adjusted accordingly before release of data and therefore not considered in the current study.

### Assessment of Risk Factors

Participants of the Framingham cohort are routinely followed up, permitting access to clinical phenotype data at multiple time points. We used first patient visit data for the offspring population, because at the time of the study only 1 visit was available for this generation.

For the parental population of the FHS, analyses were performed on values from visit 3 only, given that it had the largest number of individuals with available lipid data. The other phenotypes relevant to our study were age, body mass index (BMI), smoking status, use of lipid-lowering medications (ever treated vs. never treated), as well as the menopausal status in females. These phenotypes were used as covariates for both the offspring and parental populations. For the offspring population, we used the first adult patient visit, providing a direct adult to adult comparison.

In SardiNIA, we analyzed the same phenotypes and covariates from visit 1 in the parental and offspring population, given that both provided the largest number of patients with available lipid data.

### Statistical Analysis

#### POE on offspring lipid traits


We examined the POE on the variation of fasting lipids in the offspring populations. To identify transmission effects, we performed a series of nested, sex-stratified linear regression analyses, modeling lipid traits in offspring. The models were generated by sequentially changing the variables included; namely, all offspring covariates and corresponding parental lipid traits (Figure – Panel A). We report the adjusted  $R^2$  values throughout, which represent the proportion of variation explained by each model with all variables included in a given model.

The models assessed, also reported in Figure – Panel A, were the following:

$$\begin{aligned} \text{Model 1: Offspring Lipid Trait} \\ = \beta_0 + \beta_1(\text{Corresponding Parental Lipid Trait}) \end{aligned}$$

$$\begin{aligned} \text{Model 2: Offspring Lipid Trait} \\ = \beta_0 + \beta_1(\text{Offspring Covariate}_1) + \dots \\ + \beta_n(\text{Offspring Covariate}_n) \end{aligned}$$

$$\begin{aligned} \text{Model 3: Offspring Lipid Trait} \\ = \beta_0 + \beta_1(\text{Offspring Covariate}_1) + \dots \\ + \beta_n(\text{Offspring Covariate}_n) \\ + \beta_{n+1}(\text{Corresponding Parental Lipid Trait}) \end{aligned}$$



**Figure.** Flowchart of nested models used to determine parent of origin effects on offspring lipid traits. Parent of origin effects and relevant covariates (A) and parent of origin effects combined with SNPs previously associated with lipid traits (B). SNPs indicate single-nucleotide polymorphisms.

The summary of all analyses modeling offspring lipid traits, including the assessment of the effect of parental covariates, is provided in the Supplemental Materials (Table S1).

To evaluate the performance of each model, we compared the adjusted  $R^2$  values for each lipid trait model using a likelihood ratio test. Pair-wise model comparisons were carried out for nested pairs, namely, Model 3 versus Model 1 (effect of offspring covariates) and Model 3 versus Model 2 (effect of parental lipid trait).

### Estimating environmental effects

To estimate the effects of shared environment on lipid traits, we modeled each maternal lipid trait with the corresponding paternal lipid trait under the assumption that shared environment would be revealed by large  $R^2$  in this regression model (Model 4).

The following model was used for all parents in the trio families and then separately stratified by the sex of their offspring, to mirror the analysis above:

$$\text{Model 4: Maternal Lipid Trait} = \beta_0 + \beta_1(\text{Corresponding Paternal Lipid Trait})$$

Additional models for the effects of maternal and paternal covariates on the adjusted  $R^2$  produced by Model 4 are reported in Table S2.

To assess the role of early environment on lipid profiles, we compared sibling lipids in families with more than 1 offspring. Specifically, we determined the variance explained in same sex versus different sex sibling pairs.

### Genetic contribution to POE

To examine whether the effects of parental lipid traits are explained by genetic variants in offspring, we analyzed the effects of the 95 previously validated single-nucleotide variants (SNPs) from Teslovich et al.<sup>11</sup> on the corresponding lipid levels.

To assess the dependence of offspring lipid traits on SNPs previously associated with each lipid trait, we performed nested, sex-stratified linear regression models and compared them to the variance explained only by the corresponding parental lipid traits (Figure – Panel B).

In the Framingham cohort, genotyping was performed using the 500K Affymetrix Genechip, and many of the Teslovich SNPs were not included. We therefore used proxy variants based on high linkage disequilibrium (LD) in European populations (CEU and TSI) from phase 3 of the International HapMap Project.<sup>31–33</sup> For each nongenotyped SNP, we chose a variant on the same chromosome in strong LD ( $r^2 > 0.75$ ), having the highest minor allele frequency (list of SNPs used can be found in Table S3). Only 63 of the 95 SNPs were available either through direct genotyping or as proxies.

In contrast, in the SardiNIA cohort, genotyping information was available from 4 different Illumina arrays, one of which, Cardio-MetaboChip, included the majority of the Teslovich SNPs (Pistis et al.<sup>34</sup>). Overall, in the SardiNIA cohort, 92 of the 95 SNPs reported in Teslovich et al. were available and analyzed (Table S3). In addition, to make all analyses directly comparable between the 2 cohorts, we also evaluated the same subset of 63 SNPs, original and proxies, as in the Framingham cohort. To further assess the effects of using proxies, as opposed to the original Teslovich SNPs, in SardiNIA we also considered models using only the original 63 Teslovich SNPs for which we had either direct genotype data or proxies available in the Framingham cohort (Table S3).

We only used the subset of SNPs previously associated with each lipid trait phenotype (Table S3).

$$\text{Model 5: Offspring Lipid Trait} = \beta_0 + \beta_1(\text{Tesl. SNP}_1) + \dots + \beta_n(\text{Tesl. SNP}_n)$$

$$\text{Model 1: Offspring Lipid Trait} = \beta_0 + \beta_1(\text{Corresponding Parental Lipid Trait})$$

$$\text{Model 6: Offspring Lipid Trait} = \beta_0 + \beta_1(\text{Tesl. SNP}_1) + \dots + \beta_n(\text{Tesl. SNP}_n) + \beta_{n+1}(\text{Corresponding Parental Lipid Trait})$$

Because genotypes were not available for all participants, the number of observations in Model 1 included in this

comparison differs from the one above (Tables 7 and 8 vs. Tables 3 and 4, respectively). Analyses adding offspring covariates and using the other SNP sets for the SardiNIA cohort were also performed.

Effect sizes and potential redundancy among the influence of known genes and parental lipid trait measures were evaluated by comparing the results of likelihood ratio tests of Model 6 versus Model 1 (effect of offspring SNPs) and Model 6 versus Model 5 (effect of parental lipid trait).

All analyses were conducted using STATA (11.1; StataCorp LP, College Station, TX) and R software (R Foundation for Statistical Computing, Vienna, Austria). Two-sided *P* values are reported throughout.

## Results

### Population Characteristics

Summary statistics of lipid traits and covariates for participants are listed in Tables 1, 2, and S4, S5.

Of note, in the FHS, lipid trait measures from the offspring generation were ascertained at a younger age (mean 39.0 for daughters, 39.4 for sons) than those for parents (mean 46.4 for mothers, 48.7 for fathers). Comparisons between generations show that TC and LDL-C levels were higher in fathers than sons and higher in mothers than daughters ( $P<0.001$  for all), whereas HDL was higher in sons than fathers ( $P<0.001$ ). All comparisons are presented in Tables 1 and S4. Comparisons within each generation showed that males have higher TC, TG, and LDL, but lower HDL levels, than women. The difference between sex was statistically significant for all traits ( $P<0.003$ ), except for TC in the parents ( $P=0.06$ ).

In the SardiNIA cohort, lipid trait measures from offspring were ascertained at a younger age (mean 29.3 for daughters, 28.8 for sons) than those for their parents (mean 55.9 for mothers, 58.4 for fathers) (Table 2). Comparisons between generations showed that TC, TG, and LDL levels were higher in fathers than sons and higher in mothers than daughters ( $P<0.002$  for all). HDL was significantly higher in fathers compared to sons ( $P<0.001$ ) (Tables 1 and S5). Similarly, mothers had higher HDL compared to daughters, although not significantly. Comparisons within generations in the SardiNIA cohort showed that TG were higher in fathers than mothers ( $P<0.001$ ); HDL was significantly higher in mothers compared to fathers, as well as daughters compared to sons ( $P<0.001$  for both).

### Parent of Origin Effects on Offspring Lipid Traits

In the FHS, models including only parental lipid traits (Model 1) explained between  $\approx 1\%$  and 5% of offspring variability in 7 trait combinations and more than 5% in 9 others. The highest proportion of explained variance was  $\approx 10\%$  for mother-son HDL (Table 3; Model 1). Maternal lipids explained at least 5% of the variation in TC, LDL, and HDL of the offspring for both sex. In general, maternal lipid values provided more information regarding offspring values than did paternal values (Table 3).

For comparison, the variation of offspring lipid traits explained by all offspring covariates was more than 5% in all 16 models. The highest proportion of variability explained was  $\approx 15\%$  for LDL of daughters (Model 2). When both parental lipid traits and offspring covariates were used in a single model, the explained variation ranged from  $\approx 9\%$  (father-son LDL) to  $\approx 19\%$  (father-son HDL) (Model 3). Importantly, adding

**Table 1.** Population Characteristics of the Framingham Heart Study Participants

Lipid Traits and Risk Factors	Generation 2				Generation 3			
	n	Females*	n	Males*	n	Females*	n	Males*
Total cholesterol	416	208.81 (41.06)	416	213.76 (34.47)	228	182.73 (30.46)	188	192.29 (35.46)
Triglycerides	416	91.32 (52.57)	416	124.55 (66.49)	228	89.07 (43.51)	188	121.40 (65.55)
LDL	413	132.43 (38.34)	416	144.23 (66.49)	228	105.19 (29.27)	188	119.43 (32.40)
HDL	413	58.28 (14.15)	416	44.62 (10.39)	228	59.73 (14.38)	188	48.58 (13.02)
Age	416	46.45 (7.48)	416	48.72 (7.67)	228	39.05 (7.45)	188	39.39 (7.56)
BMI	413	24.83 (4.93)	414	27.14 (3.30)	226	25.52 (5.58)	188	27.97 (4.71)
Anticholesterol treatment	416	5 (1.20)	416	7 (1.68)	228	5 (2.19)	188	17 (9.04)
Smoking status	415	96 (23.13)	416	91 (21.88)	228	94 (41.23)	188	61 (32.45)
Menopause	416	159 (38.22)			228	24 (10.53)		

BMI indicates body mass index; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

\*Mean (SD) for continuous variables, n (% total) for categorical variables.



**Table 2.** Population Characteristics of the SardiNIA Cohort

Lipid Traits and Risk Factors	Generation 2				Generation 3			
	n	Females*	n	Males*	n	Females*	n	Males*
Total cholesterol	277	221.90 (39.40)	151	223.13 (41.56)	168	194.43 (37.83)	136	188.54 (46.68)
Triglycerides	277	85.26 (50.44)	147	105.57 (60.87)	168	71.92 (40.14)	135	82.17 (49.91)
LDL	277	136.51 (34.28)	147	140.76 (34.98)	168	113.41 (29.86)	135	116.55 (37.99)
HDL	277	68.34 (16.05)	151	59.38 (13.23)	168	66.63 (16.29)	136	54.49 (11.83)
Age	277	55.94 (11.41)	151	58.40 (10.71)	168	29.26 (10.41)	136	28.80 (9.67)
BMI	277	27.22 (5.03)	151	27.98 (3.88)	168	22.28 (3.28)	136	24.49 (3.89)
Anticholesterol treatment	277	0	151	1 (0.06)	168	0	136	0
Smoking status	277	29 (10.47)	151	44 (29.14)	168	34 (20.24)	136	51 (37.50)
Menopause	277	175 (63.18)			168	9 (5.36)		

BMI indicates body mass index; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

\*Mean (SD) for continuous variables, n (% total) for categorical variables.

parental lipid trait values to a model containing offspring covariates significantly increased the amount of variation explained for TC, LDL, and HDL in all parent offspring pairs (Table 3, *P* values M3/M2).

The amount of variation explained in TG models for fathers-daughters was significant, whereas all other TG parent-offspring pairs were below the significance level. This indicates that the effects of parental lipid traits and offspring

covariates are significant and independent of one another, with the exception of TG measures. Parental covariates explained less of offspring lipid variability than either offspring covariates or parental lipids as can be seen by comparison of models 2, 3, S1, and S2 (Table S6).

In SardiNIA, models of offspring lipid traits using only corresponding parental lipid traits explained between 0.01% and 5% of in 6 models and more than 5% in 9 others. One

**Table 3.** Estimating the Parent of Origin Effects on Lipid Traits in the Framingham Heart Study

Parent-Offspring Pair	Modeled Lipid Trait	Adjusted $R^2$			Likelihood Ratio Tests	
		Model 1	Model 2	Model 3	<i>P</i> Value M3/M1	<i>P</i> Value M3/M2
Mothers daughters	TC (n=223)	0.0789	0.1188	0.1469	<0.001	0.004
	TG (n=223)	0.022	0.1137	0.1237	<0.001	0.059
	LDL (n=222)	0.0808	0.1307	0.1723	<0.001	<0.001
	HDL (n=222)	0.0691	0.1183	0.167	<0.001	<0.001
Fathers-daughters	TC (n=224)	0.0335	0.1314	0.1472	<0.001	0.024
	TG (n=224)	0.0207	0.1163	0.1295	<0.001	0.036
	LDL (n=224)	0.0547	0.1469	0.1722	<0.001	0.005
	HDL (n=224)	0.0762	0.1147	0.1648	<0.001	<0.001
Mothers-sons	TC (n=187)	0.0552	0.0676	0.114	0.004	0.001
	TG (n=187)	0.0182	0.1066	0.1175	<0.001	0.068
	LDL (n=185)	0.0588	0.0661	0.1247	0.002	<0.001
	HDL (n=185)	0.0963	0.1291	0.1808	<0.001	<0.001
Fathers-sons	TC (n=188)	0.0172	0.0688	0.0932	<0.001	0.014
	TG (n=188)	0.029	0.11	0.1206	<0.001	0.070
	LDL (n=188)	0.0098	0.0696	0.0884	<0.001	0.027
	HDL (n=188)	0.0784	0.1307	0.1862	<0.001	<0.001

HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.

model explained less than 0.01% of the variability (TG for fathers-daughters). The highest proportion of variability explained was  $\approx 15\%$  for mother-son LDL (Table 4; Model 1). As in the FHS, maternal lipid traits in SardinIA explained at least 5% of the corresponding variation for TC, LDL, and HDL of both sons and daughters and generally performed better than paternal models.

The variation of offspring lipid traits explained by offspring covariates alone was between 0.01% and 5% in 4 models and more than 5% in 12 models (Model 2). The highest proportion of variability explained was  $\approx 36\%$  for TC of sons (Model 2). When both parental lipid traits and offspring covariates were used in the same model, 1 model explained marginal variability ( $<0.01\%$ ), 1 model explained between 0.01% and 5%, and 14 models explained more than 5%. The highest proportion of variability explained was  $\approx 39\%$  for father-son TC (Model 3). Adding parental lipid trait values to a model containing offspring covariates explained TC significantly better when modeling sons' levels with fathers' or mothers' (Table 4, *P* values M3/M2), whereas results for daughters trended in the same direction. With the exception of fathers-sons, adding parental HDL or LDL values to Model 2 significantly improved all parent-offspring pair models (Table 4).

Maternal TG levels explained significantly more of daughters' TG, but had no effect on the other parent-offspring pairs (Table 4). As with the Framingham results, parental covariates generally explained less of the offspring lipid variability than either offspring covariates or parental lipids, as can be seen by comparison of models 2, 3, S1, and S2 (Table S7).

## Environmental Effects

In the FHS, the variation of maternal lipid traits explained by the corresponding paternal lipid traits ranged from negligible ( $<0.01\%$ ) for TG to  $\approx 1\%$  for TC and LDL (Table 5). In SardinIA, the percentage of maternal lipid traits explained by corresponding paternal lipid traits ranged from negligible ( $<0.01\%$ ) for TG to  $\approx 1\%$  for TC, LDL, and HDL (Table 6; Supplementary Results). Both results indicate that shared adult environments do not significantly impact our findings.

In the SardinIA cohort, when only parents of daughters were considered, paternal lipid traits explained up to 2.5% of variation for HDL. When only parents of sons were considered, paternal lipid traits explained up to 2.5% of LDL variation (Table 6). These results indicate that the shared environment of parents explains very little of lipid trait variation in the unrelated parent pairs, and suggest that the effects of

**Table 4.** Estimating the Parent of Origin Effects on Lipid Traits in the SardinIA Cohort

Parent-Offspring Pair	Modeled Lipid Trait	Adjusted $R^2$			Likelihood Ratio Tests	
		Model 1	Model 2	Model 3	<i>P</i> Value M3/M1	<i>P</i> Value M3/M2
Mothers-daughters	TC (n=152)	0.051	0.159	0.171	$<0.001$	0.071
	TG (n=152)	0.045	0.107	0.126	0.002	0.038
	LDL (n=152)	0.067	0.129	0.164	$<0.001$	0.007
	HDL (n=152)	0.102	0.053	0.157	0.008	$<0.001$
Fathers-daughters	TC (n=86)	0.020	0.066	0.087	0.035	0.083
	TG (n=84)	$<0.001$	0.002	$<0.001$	0.358	0.866
	LDL (n=84)	0.114	0.030	0.193	0.017	$<0.001$
	HDL (n=86)	0.078	0.062	0.121	0.079	0.010
Mothers-sons	TC (n=125)	0.145	0.312	0.362	$<0.001$	0.001
	TG (n=125)	0.024	0.188	0.185	$<0.001$	0.442
	LDL (n=125)	0.147	0.308	0.385	$<0.001$	$<0.001$
	HDL (n=125)	0.056	0.045	0.079	0.107	0.019
Fathers-sons	TC (n=65)	0.066	0.361	0.388	$<0.001$	0.050
	TG (n=63)	0.008	0.123	0.130	0.010	0.218
	LDL (n=63)	0.007	0.277	0.289	$<0.001$	0.144
	HDL (n=65)	0.037	0.010	0.032	0.414	0.111

The overall numbers of parents used for this analysis are lower than the numbers used in estimating parent of origin effects in Table 2 because, unlike the Framingham Heart Study, the SardinIA cohort is comprised of more single-parent families and, consequently, fewer complete trios. HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.



**Table 5.** Estimating the Effects of Shared Environment in the Framingham Heart Study

Modeled Maternal Lipid Trait	Adjusted $R^2$ From Modeling With Corresponding Paternal Lipid Trait					
	n	All	n	Parents of Daughters Only	n	Parents of Sons Only
TC	416	0.014	223	0.027	187	<0.001
TG	416	<0.001	223	<0.001	187	<0.001
LDL	413	0.012	222	0.032	185	<0.001
HDL	413	0.005	222	0.004	185	0.001

HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.

parental lipids on offspring are linked to factors unrelated to the shared environment.

When only parents of daughters were considered, paternal lipid traits explained between 0.02% and 3% of maternal TG and LDL, respectively. When only parents of sons were considered, paternal lipid traits only explained  $\approx 0.1\%$  of maternal HDL variation, with negligible variation explained for all other lipid traits (Table 5). Using maternal lipid traits to model corresponding paternal lipid traits, produced similar results to using paternal lipid traits to model maternal traits when covariates were included (Tables S8 and S9).

The role of shared early environment was also assessed by comparing variance explained for each lipid trait values in siblings of the same to those of the opposite sex. Variance explained between siblings of the same sex ranged between 5% and 12% for females and between 0.5% and 6% for males, whereas between offspring of opposite sex the results were generally smaller (0% to 5%). This supports the conclusion that the results between parents and offspring are not attributable to shared environment (Table S10).

**Table 6.** Estimating the Effects of Shared Environment in the SardiNIA Cohort

Modeled Maternal Lipid Trait	Adjusted $R^2$ From Modeling With Corresponding Paternal Lipid Trait					
	n	All	n	Parents of Daughters Only	n	Parents of Sons Only
TC	126	0.014	72	0.003	54	0.013
TG	123	<0.001	70	<0.001	53	<0.001
LDL	123	0.010	70	<0.001	53	0.025
HDL	126	0.012	72	0.025	54	<0.001

HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.

## Genetic Contribution to the POE

In the FHS, SNPs previously associated with each lipid trait explained a negligible amount of variability (<0.01%) in 2 models, between 0.01% and 5% in 8 models, and more than 5% in 6 models, the highest being  $\approx 10\%$  for daughters' TG (Table 7; Model 5). In the offspring with available genotype data (a subset of individuals from Model 1), parental lipid traits explained between 0.01% and 5% of variability in 8 models and more than 5% in 8 models, the highest being  $\approx 10\%$  for sons' HDL with paternal measures (Table 7; Model 1). Adding parental lipid values to the model containing all SNPs produced significantly better models in explaining TC, LDL, and HDL in all parent-offspring pairs, except fathers-sons, where only the HDL model was significantly improved (Table 7,  $P$  values M6/M5). Conversely, adding all SNPs to models containing parental lipid values significantly improved HDL in the mothers-sons model and all models of TGs, with the exception of the fathers-sons comparison (Table 7,  $P$  values M6/M1).

In SardiNIA, models with the 92 SNPs previously associated with lipid traits (Table 8) resulted in negligible percent variance explained (<0.01%) in 8 of the 16 models (Table 8; Model 5). Four models explained between 0.01% and 5%, and 4 explained more than 5%, with the highest percentage being  $\approx 30\%$  for HDL in mother-son pairs (Model 5). In the subset of offspring with available genotype data, the variation of offspring measures explained by parental lipid traits only was negligible for 2 models (sons' and daughters' TG with paternal levels) (Model 1). Parental lipid traits explained between 0.01% and 5% in 4 models, and greater than 5% in 10 models, the highest being  $\approx 16\%$  for sons' TC with maternal levels (Table 8; Model 1). Adding parental lipid values to the model containing all SNPs significantly improved the models for TC, LDL, and HDL in mother-daughter and father-daughter models, for all lipid traits in mother-son models, and for TC and HDL in father-son models (Table 8,  $P$  values M6/M5). Adding all SNPs to a model containing parental lipid values produced significant results for HDL in mother-daughter models, for TC and HDL in father-daughter models, for HDL in mother-son models, and for TC, TG, and HDL in father-son models (Table 8,  $P$  values M6/M1). These results were generally consistent when the alternative SNP sets were used; namely, the subset of 63 original and proxy SNPs from the Framingham analysis, and also when all nonproxy 63 SNPs (Tables S11 through S16).

## Discussion

We investigated how parental serum levels of TC, TG, LDL, and HDL can be used to model lipid traits of the offspring, using sex-stratified analyses. We also compared the effect of

**Table 7.** Comparing Lipid Trait Relevant SNPs to Parent of Origin Effects in the Framingham Heart Study

Parent-Offspring Pair	Modeled Lipid Trait	Adjusted $R^2$			Likelihood Ratio Tests	
		Model 5	Model 1	Model 6	P Value M6/M5	P Value M6/M1
Mothers-daughters	TC (n=186)	0.005	0.079	0.089	<0.001	0.237
	TG (n=186)	0.102	0.023	0.111	0.091	0.008
	LDL (n=185)	0.024	0.091	0.125	<0.001	0.094
	HDL (n=184)	<0.001	0.095	0.107	<0.001	0.216
Fathers-daughters	TC (n=186)	0.005	0.015	0.023	0.033	0.261
	TG (n=186)	0.102	0.017	0.105	0.182	0.008
	LDL (n=186)	0.025	0.036	0.077	0.001	0.076
	HDL (n=185)	<0.001	0.101	0.068	<0.001	0.101
Mothers-sons	TC (n=153)	0.039	0.064	0.079	0.006	0.199
	TG (n=153)	0.079	0.013	0.074	0.607	0.050
	LDL (n=152)	0.003	0.068	0.064	0.001	0.404
	HDL (n=152)	0.061	0.091	0.166	<0.001	0.025
Fathers-sons	TC (n=153)	0.040	0.004	0.040	0.251	0.116
	TG (n=153)	0.079	0.020	0.080	0.249	0.052
	LDL (n=153)	0.007	0.003	0.019	0.087	0.236
	HDL (n=151)	0.064	0.104	0.155	<0.001	0.059

For models 5 and 6, there are 4 original, 25 proxy for TC; 4 original, 17 proxy for TG; 3 original, 15 proxy for LDL; 5 original, 25 proxy for HDL. HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; SNP, single-nucleotide polymorphism; TC, total cholesterol; TG, triglycerides.

**Table 8.** Comparing Lipid Trait Relevant SNPs to Parent of Origin Effects in the SardinIA Cohort

Parent-Offspring Pair	Modeled Lipid Trait	Adjusted $R^2$			Likelihood Ratio Tests	
		Model 5	Model 1	Model 6	P Value M6/M5	P Value M6/M1
Mothers-daughters	TC (n=133)	0.037	0.055	0.069	0.015	0.100
	TG (n=133)	<0.001	0.046	<0.001	0.063	0.740
	LDL (n=133)	0.061	0.081	0.102	0.007	0.158
	HDL (n=133)	0.041	0.072	0.158	<0.001	0.014
Fathers-daughters	TC (n=78)	0.054	0.007	0.153	0.001	0.001
	TG (n=76)	<0.001	<0.001	<0.001	0.686	0.455
	LDL (n=76)	<0.001	0.095	0.068	0.008	0.168
	HDL (n=78)	0.003	0.077	0.163	<0.001	0.003
Mothers-sons	TC (n=109)	<0.001	0.169	0.059	<0.001	0.494
	TG (n=109)	<0.001	0.032	0.022	0.038	0.240
	LDL (n=109)	<0.001	0.152	0.119	<0.001	0.403
	HDL (n=109)	0.304	0.069	0.353	0.002	<0.001
Fathers-sons	TC (n=60)	<0.001	0.062	<0.001	0.001	0.006
	TG (n=59)	0.151	<0.001	0.138	0.253	0.004
	LDL (n=59)	<0.001	<0.001	<0.001	0.383	0.681
	HDL (n=60)	0.009	0.057	0.198	<0.001	<0.001

For models 5 and 6, there are 51 SNPs for TC, 32 SNPs for TG, 37 SNPs for LDL, and 47 SNPs for HDL. HDL indicates high-density lipoprotein; LDL, low-density lipoprotein; SNP, single-nucleotide polymorphism; TC, total cholesterol; TG, triglycerides.

parental lipids with that of known and validated genetic variants in loci previously associated with plasma lipid levels.<sup>11</sup> These analyses were performed in 2 well-characterized and large prospective cohorts, the FHS (inclusive of Offspring and Generation 3 cohorts) and the SardiNIA Study.

The most important finding from our study is that, in general, parental serum lipid levels explain a higher proportion of variability in the offspring than do the 95 loci described in the work by Teslovich et al.<sup>11</sup> This effect was not owing to shared environment and was independent of nongenetic factors known to modulate lipid levels. These results were generally consistent between the Framingham and SardiNIA cohorts (Tables 3 through 8). Given the independence from other variables and from the currently known genetic loci, these results suggest the presence of unknown variants or mechanisms responsible for the missing heritability of lipid traits<sup>12,13</sup> and serve to emphasize the size of the gap in our knowledge of factors that affect lipid levels. However, because we do not yet have an understanding of these other determinants, we argue that parental lipid levels explain those of adult offspring better than do the validated variants in 95 genes. Thus, knowledge of parental lipid levels provides information to predict future lipid levels in the offspring and should be used as a tool to target pediatric lipid testing.

With the exception of Mendelian forms of dyslipidemia, serum lipids are complex traits influenced by multiple genetic and nongenetic factors. Therefore, the use of single-gene variants is of little utility in the prediction of this complex phenotype. To date, GWA studies have identified risk loci that have high statistical significance, but low biological effect sizes, and such markers are not generally practical for predictive purposes.<sup>35–38</sup> As a consequence, genetics-based predictions using multilocus modeling thus far provide marginal clinical utility in prevention because they have low predictive power.<sup>35,39</sup> The value of a genetic test depends on several factors, including the number of genes influencing the trait, frequency of the associating allele, and strength of association between genotype and phenotype, making accurate predictions from simple models extremely difficult.<sup>35</sup> In contrast, the results of our study show that family history and nongenetic covariates better explain lipid levels in adult offspring than does variation in the loci known to influence lipids across study populations. Our findings are in agreement with what has been shown in other complex phenotypes, such as type II diabetes, where risk scores not including genetic variant data were virtually identical to those incorporating validated genes for type II diabetes risk.<sup>37,38</sup> Similar results were also found in a previous study where a gene-based score did not significantly improve the association between canonical risk factors and CVD.<sup>39</sup> Recently, 62 additional lipid-associated loci were identified, but their effects were small, explaining <2% of the total phenotype variance and therefore

should not substantially impact our conclusions.<sup>40</sup> Furthermore, we demonstrate the existence of parent-of-origin effects on lipid levels, which are sex-specific and likely owing to both genetic and epigenetic factors. Such effects have been shown to modulate some of the risk factors for dyslipidemia. For example, a recent GWA study has demonstrated parent-of-origin effects in the degree to which SNPs in 2 genes, *SLC2A10* and *KCNK9*, affect BMI, a major factor affecting lipids. These SNPs showed “polar overdominance,” where homozygotes of either SNP had the same average BMI, whereas heterozygotes differed as a function of parent of origin.<sup>41</sup>

We also found that maternal traits generally explain more of the offspring's TC, HDL, and LDL. Maternal lipid traits explained at least 5% of the offspring variability in TC, HDL, and LDL of both sons and daughters in both cohorts. Paternal traits were less consistent, given that their effects ranged from nonsignificant in multiple traits to relatively high in explaining the daughter's LDL and HDL (5% to 8% of explained variability in Framingham and 7% to 11% in SardiNIA). The finding of stronger maternal influences on offspring lipid traits is consistent with epidemiological data demonstrating that maternal lifestyle and environment (such as nutritional status, stress level, insulin resistance, diabetes, hypertension, hypercholesterolemia, obesity, and smoking), both at time of conception and during pregnancy, influence the offspring's phenotypes, such as adiposity, blood pressure, fatty streak formation, or diabetes.<sup>15,17,42–50</sup> This is consistent with the Barker hypothesis, that is, that early exposure, both pre- and postnatal, can affect risk of adult-onset disease.<sup>51,52</sup>

Interestingly, small or nonexistent parent-of-origin effects were generally observed for TG (Tables 3 and 4). TGs also provided the most variable results when using genetic models (Tables 5 and 6). TG levels have a smaller parent-of-origin effect than the other lipid traits, and a bigger part of their heritability may be determined by other factors, such as rare variants. It has been recently demonstrated that rare APOC3 mutations have a strong influence on plasma TG levels in aggregate.<sup>53,54</sup>

Our study has several limitations. Although we were able to identify models that account for a significant portion of lipid variation explained, we were not able to provide a mechanism for this effect. We can only speculate that our observations may forecast discovery of additional genes, gene-gene interactions, or epigenetic effects that regulate lipid levels. Furthermore, in our analyses, we did not account for specific environmental variables, such as diet, alcohol, exercise, socioeconomic status, and use of specific medications. However, it is of note that the 2 cohorts we studied would be expected to have different environmental exposures and the results were still mostly concordant. This discrepancy is not likely to have influenced the results, and it would have had

an attenuating effect even if it did. Both cohorts are prospective studies analyzing populations of European ancestry, but Framingham's residents are from multiple European origins, whereas the participants in SardinIA are part of a genetic isolate. This may be the basis for the minor discrepancies we observed (seen in Tables 3 through 6). Dietary habits were not directly quantified and, consequently, were not represented in our models in either cohort, although BMI and lipid medication covariates can be considered partial proxies for diet and lifestyle. However, the similarities of results between cohorts provide additional strength to our main claim.

In conclusion, we have determined that parent-of-origin effects explain more variability in the adult offspring's lipid levels than do common variants in the loci known to modulate lipid metabolism. Knowledge of the parent's lipid levels may provide an inexpensive and practical means to predict future lipid levels in their children.

## Acknowledgments

**Author Contributions:** Analyses were performed by Predazzi (Vanderbilt University, now at Oregon Health and Science University), Sobota (Vanderbilt University, Dartmouth College), Bartlett (Dartmouth College) and Sanna (Istituto di Ricerca Genetica e Biomedica del Consiglio Nazionale delle Ricerche). Predazzi and Sanna had full access to the data in the study at the time in which the analyses were performed and take full responsibility for the credibility and the accuracy of the data analysis.

## Sources of Funding

This work was partially funded by grant NIH 2R01HL057986-15A1 and 5R01HL106845-03 (Fazio), NIH 2T32HL007751-16A2 (Predazzi), NIH P20 GM103534 (Williams), NIH HL116263 (Linton), NIH National Institute of Aging N01-AG-1-2109, N01-AG-1-2109, and Sardinian Autonomous Region (L.R. no. 7/2009) cRP3-154 (Cucca).

Sponsors/funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; as well as in the preparation, review, or approval of the manuscript.

## Disclosures

None.

## References

- Kathiresan S, Manning AK, Demissie S, D'Agostino RB, Surti A, Guiducci C, Gianniny L, Burt NP, Melander O, Orho-Melander M, Arnett DK, Peloso GM, Ordovas JM, Cupples LA. A genome-wide association study for blood lipid

phenotypes in the Framingham Heart Study. *BMC Med Genet*. 2007;8(suppl 1):S17.

- Hunt SC, Hasstedt SJ, Kuida H, Stults BM, Hopkins PN, Williams RR. Genetic heritability and common environmental components of resting and stressed blood pressures, lipids, and body mass index in Utah pedigrees and twins. *Am J Epidemiol*. 1989;129:625–638.
- Weiss LA, Pan L, Abney M, Ober C. The sex-specific genetic architecture of quantitative traits in humans. *Nat Genet*. 2006;38:218–222.
- Diabetes Genetics Initiative of Broad Institute of H, Mit LU, Novartis Institutes of BioMedical R, Saxena R, Voight BF, Lyssenko V, Burt NP, de Bakker PI, Chen H, Roix JJ, Kathiresan S, Hirschhorn JN, Daly MJ, Hughes TE, Groop L, Altshuler D, Almgren P, Florez JC, Meyer J, Ardlie K, Bengtsson Bostrom K, Isomaa B, Lettre G, Lindblad U, Lyon HN, Melander O, Newton-Cheh C, Nilsson P, Orho-Melander M, Rastam L, Speliotes EK, Taskinen MR, Tuomi T, Guiducci C, Berglund A, Carlson J, Gianniny L, Hackett R, Hall L, Holmkvist J, Laurila E, Sjogren M, Sterner M, Surti A, Svensson M, Svensson M, Tewhey R, Blumenstiel B, Parkin M, Defelice M, Barry R, Brodeur W, Camarata J, Chia N, Fava M, Gibbons J, Handsaker B, Healy C, Nguyen K, Gates C, Sougnez C, Gage D, Nizzari M, Gabriel SB, Chirn GW, Ma Q, Parikh H, Richardson D, Ricke D, Purcell S. Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science*. 2007;316:1331–1336.
- Willer CJ, Sanna S, Jackson AU, Scuteri A, Bonnycastle LL, Clarke R, Heath SC, Timpson NJ, Najjar SS, Stringham HM, Strait J, Duren WL, Maschio A, Busonero F, Mulas A, Albai G, Swift AJ, Morken MA, Narisu N, Bennett D, Parish S, Shen H, Galan P, Meneton P, Hercberg S, Zelenika D, Chen WM, Li Y, Scott LJ, Scheet PA, Sundvall J, Watanabe RM, Nagaraja R, Ebrahim S, Lawlor DA, Ben-Shlomo Y, Davey-Smith G, Shuldiner AR, Collins R, Bergman RN, Uda M, Tuomilehto J, Cao A, Collins FS, Lakatta E, Lathrop GM, Boehnke M, Schlessinger D, Mohlke KL, Abecasis GR. Newly identified loci that influence lipid concentrations and risk of coronary artery disease. *Nat Genet*. 2008;40:161–169.
- Kathiresan S, Melander O, Guiducci C, Surti A, Burt NP, Rieder MJ, Cooper GM, Roos C, Voight BF, Havulinna AS, Wahlstrand B, Hedner T, Corella D, Tai ES, Ordovas JM, Berglund G, Vartiainen E, Jousilahti P, Hedblad B, Taskinen MR, Newton-Cheh C, Salomaa V, Peltonen L, Groop L, Altshuler DM, Orho-Melander M. Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. *Nat Genet*. 2008;40:189–197.
- Aulchenko YS, Ripatti S, Lindqvist I, Boomsma D, Heid IM, Pramstaller PP, Penninx BW, Janssens AC, Wilson JF, Spector T, Martin NG, Pedersen NL, Kyvik KO, Kaprio J, Hofman A, Freimer NB, Jarvelin MR, Gyllenstein U, Campbell H, Rudan I, Johansson A, Marroni F, Hayward C, Vitart V, Jonasson I, Pattaro C, Wright A, Hastie N, Pichler I, Hicks AA, Falchi M, Willemsen G, Hottenga JJ, de Geus EJ, Montgomery GW, Whitfield J, Magnusson P, Saharinen J, Perola M, Silander K, Isaacs A, Sijbrands EJ, Uitterlinden AG, Witteman JC, Oostra BA, Elliott P, Ruokonen A, Sabatti C, Gieger C, Meitinger T, Kronenberg F, Doring A, Wichmann HE, Smit JH, McCarthy MI, van Duijn CM, Peltonen L; Consortium E. Loci influencing lipid levels and coronary heart disease risk in 16 European population cohorts. *Nat Genet*. 2009;41:47–55.
- Wallace C, Newhouse SJ, Braund P, Zhang F, Tobin M, Falchi M, Ahmadi K, Dobson RJ, Marciano AC, Hajat C, Burton P, Deloukas P, Brown M, Connell JM, Dominiczak A, Lathrop GM, Webster J, Farrall M, Spector T, Samani NJ, Caulfield MJ, Munroe PB. Genome-wide association study identifies genes for biomarkers of cardiovascular disease: serum urate and dyslipidemia. *Am J Hum Genet*. 2008;82:139–149.
- Sabatti C, Service SK, Hartikainen AL, Pouta A, Ripatti S, Brodsky J, Jones CG, Zaitlen NA, Varilo T, Kaakinen M, Sovio U, Ruokonen A, Laitinen J, Jakkula E, Coin L, Hoggart C, Collins A, Turunen H, Gabriel S, Elliot P, McCarthy MI, Daly MJ, Jarvelin MR, Freimer NB, Peltonen L. Genome-wide association analysis of metabolic traits in a birth cohort from a founder population. *Nat Genet*. 2009;41:35–46.
- Chasman DI, Pare G, Mora S, Hopewell JC, Peloso G, Clarke R, Cupples LA, Hamsten A, Kathiresan S, Malarstig A, Ordovas JM, Ripatti S, Parker AN, Miletich JP, Ridker PM. Forty-three loci associated with plasma lipoprotein size, concentration, and cholesterol content in genome-wide analysis. *PLoS Genet*. 2009;5:e1000730.
- Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, Pirruccello JP, Ripatti S, Chasman DI, Willer CJ, Johansen CT, Fouchier SW, Isaacs A, Peloso GM, Barbalic M, Ricketts SL, Bis JC, Aulchenko YS, Thorleifsson G, Feitosa MF, Chambers J, Orho-Melander M, Melander O, Johnson T, Li X, Guo X, Li M, Shin Cho Y, Jin Go M, Jin Kim Y, Lee JY, Park T, Kim K, Sim X, Tsee-Hee Ong R, Croteau-Chonka DC, Lange LA, Smith JD, Song K, Hua Zhao J, Yuan X, Luan J, Lamina C, Ziegler A, Zhang W, Zee RY, Wright AF, Witteman JC, Wilson JF, Willemsen G, Wichmann HE, Whitfield JB, Waterworth DM, Wareham NJ, Waeber G, Vollenweider P, Voight BF, Vitart V, Uitterlinden AG, Uda M, Tuomilehto J, Thompson JR, Tanaka T, Surakka I, Stringham HM, Spector TD, Soranzo N, Smit JH, Sinisalo J, Silander K, Sijbrands EJ, Scuteri A, Scott J, Schlessinger D, Sanna S, Salomaa V, Saharinen J, Sabatti C, Ruokonen A, Rudan I, Rose LM, Roberts R, Rieder M, Psaty BM, Pramstaller PP, Pichler I,



- Perola M, Penninx BW, Pedersen NL, Pattaro C, Parker AN, Pare G, Oostra BA, O'Donnell CJ, Nieminen MS, Nickerson DA, Montgomery GW, Meitinger T, McPherson R, McCarthy MI, McArdle W, Masson D, Martin NG, Marroni F, Mangino M, Magnusson PK, Lucas G, Luben R, Loos RJ, Lokki ML, Lettre G, Langenberg C, Launer LJ, Lakatta EG, Laaksonen R, Kyvik KO, Kronenberg F, Konig IR, Khaw KT, Kaprio J, Kaplan LM, Johansson A, Jarvelin MR, Janssens AC, Ingelsson E, Igl W, Kees Hovingh G, Hottenga JJ, Hofman A, Hicks AA, Hengstenberg C, Heid IM, Hayward C, Havulinna AS, Hastie ND, Harris TB, Haritunians T, Hall AS, Gyllenstein U, Guiducci C, Groop LC, Gonzalez E, Gieger C, Freimer NB, Ferrucci L, Erdmann J, Elliott P, Ejebe KG, Doring A, Dominiczak AF, Demissie S, Deloukas P, de Geus EJ, de Faire U, Crawford G, Collins FS, Chen YD, Caulfield MJ, Campbell H, Burtt NP, Bonnycastle LL, Boomsma DI, Boekholdt SM, Bergman RN, Barroso I, Bandinelli S, Ballantyne CM, Assimes TL, Quertermous T, Altschuler D, Seielstad M, Wong TY, Tai ES, Feranil AB, Kuzawa CW, Adair LS, Taylor HA Jr, Borecki IB, Gabriel SB, Wilson JG, Holm H, Thorsteinsdottir U, Gudnason V, Krauss RM, Mohlke KL, Ordovas JM, Munroe PB, Kooner JS, Tall AR, Hegele RA, Kastelein JJ, Schadt EE, Rotter JI, Boerwinkle E, Strachan DP, Mooser V, Stefansson K, Reilly MP, Samani NJ, Schunkert H, Cupples LA, Sandhu MS, Ridker PM, Rader DJ, van Duijn CM, Peltonen L, Abecasis GR, Boehnke M, Kathiresan S. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*. 2010;466:707–713.
12. Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, Nadeau JH. Missing heritability and strategies for finding the underlying causes of complex disease. *Nat Rev Genet*. 2010;11:446–450.
  13. Manolio TA. Genomewide association studies and assessment of the risk of disease. *N Engl J Med*. 2010;363:166–176.
  14. Ravelli AC, van der Meulen JH, Michels RP, Osmond C, Barker DJ, Hales CN, Bleker OP. Glucose tolerance in adults after prenatal exposure to famine. *Lancet*. 1998;351:173–177.
  15. Palinski W, Yamashita T, Freigang S, Napoli C. Developmental programming: maternal hypercholesterolemia and immunity influence susceptibility to atherosclerosis. *Nutr Rev*. 2007;65:S182–S187.
  16. Napoli C, Infante T, Casamassimi A. Maternal-foetal epigenetic interactions in the beginning of cardiovascular damage. *Cardiovasc Res*. 2011;92:367–374.
  17. Palinski W, Nicolaides E, Liguori A, Napoli C. Influence of maternal dysmetabolic conditions during pregnancy on cardiovascular disease. *J Cardiovasc Transl Res*. 2009;2:277–285.
  18. van der Graaf A, Vissers MN, Gaudet D, Brisson D, Sivapalaratnam S, Roseboom TJ, Jansen AC, Kastelein JJ, Hutten BA. Dyslipidemia of mothers with familial hypercholesterolemia deteriorates lipids in adult offspring. *Arterioscler Thromb Vasc Biol*. 2010;30:2673–2677.
  19. Rutherford JD. Maternal heterozygous familial hypercholesterolemia and its consequences for mother and child. *Circulation*. 2011;124:1599–1601.
  20. Churchill D, Perry IJ, Beevers DG. Ambulatory blood pressure in pregnancy and fetal growth. *Lancet*. 1997;349:7–10.
  21. Mott R, Yuan W, Kaisaki P, Gan X, Cleak J, Edwards A, Baud A, Flint J. The architecture of parent-of-origin effects in mice. *Cell*. 2014;156:332–342.
  22. Ideraabdullah FY, Kim K, Pomp D, Moran JL, Beier D, de Villena FP. Rescue of the mouse DDK syndrome by parent-of-origin-dependent modifiers. *Biol Reprod*. 2007;76:286–293.
  23. Eberle C, Merki E, Yamashita T, Johnson S, Armando AM, Quehenberger O, Napoli C, Palinski W. Maternal immunization affects in utero programming of insulin resistance and type 2 diabetes. *PLoS One*. 2012;7:e45361.
  24. Chadwick LH, Willard HF. Genetic and parent-of-origin influences on X chromosome choice in Xce heterozygous mice. *Mamm Genome*. 2005;16:691–699.
  25. Hines IN, Hartwell HJ, Feng Y, Theve EJ, Hall GA, Hashway S, Connolly J, Fecteau M, Fox JG, Rogers AB. Insulin resistance and metabolic hepatocarcinogenesis with parent-of-origin effects in AxB mice. *Am J Pathol*. 2011;179:2855–2865.
  26. Liljelund P, Handforth A, Homanics GE, Olsen RW. GABAA receptor beta3 subunit gene-deficient heterozygous mice show parent-of-origin and gender-related differences in beta3 subunit levels, EEG, and behavior. *Brain Res Dev Brain Res*. 2005;157:150–161.
  27. Dawber TR, Meadors GF, Moore FE Jr. Epidemiological approaches to heart disease: the Framingham Study. *Am J Public Health Nations Health*. 1951;41:279–281.
  28. Feinleib M, Kannel WB, Garrison RJ, McNamara PM, Castelli WP. The Framingham Offspring Study. Design and preliminary data. *Prev Med*. 1975;4:518–525.
  29. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
  30. Pilia G, Chen WM, Scuteri A, Orru M, Albai G, Dei M, Lai S, Usala G, Lai M, Loi P, Mameli C, Vacca L, Deiana M, Olla N, Masala M, Cao A, Najjar SS, Terracciano A, Nedorezov T, Sharov A, Zonderman AB, Abecasis GR, Costa P, Lakatta E, Schlessinger D. Heritability of cardiovascular and personality traits in 6,148 Sardinians. *PLoS Genet*. 2006;2:e132.
  31. International HapMap C. A haplotype map of the human genome. *Nature*. 2005;437:1299–1320.
  32. International HapMap C, Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, Belmont JW, Boudreau A, Hardenbol P, Leal SM, Pasternak S, Wheeler DA, Willis TD, Yu F, Yang H, Zeng C, Gao Y, Hu H, Hu W, Li C, Lin W, Liu S, Pan H, Tang X, Wang J, Wang W, Yu J, Zhang B, Zhang Q, Zhao H, Zhao H, Zhou J, Gabriel SB, Barry R, Blumenstiel B, Camargo A, Defelice M, Faggart M, Goyette M, Gupta S, Moore J, Nguyen H, Onofrio RC, Parkin M, Roy J, Stahl E, Winchester E, Ziaugra L, Altschuler D, Shen Y, Yao Z, Huang W, Chu X, He Y, Jin L, Liu Y, Shen Y, Sun W, Wang H, Wang Y, Wang Y, Xiong X, Xu L, Wayne MM, Tsui SK, Xue H, Wong JT, Galver LM, Fan JB, Gunderson K, Murray SS, Oliphant AR, Chee MS, Montpetit A, Chagnon F, Ferretti V, Leboeuf M, Olivier JF, Phillips MS, Roumy S, Sallee C, Verner A, Hudson TJ, Kwok PY, Cai D, Koboldt DC, Miller RD, Pawlikowska L, Taillon-Miller P, Xiao M, Tsui LC, Mak W, Song YQ, Tam PK, Nakamura Y, Kawaguchi T, Kitamoto T, Morizono T, Nagashima A, Ohnishi Y, Sekine A, Tanaka T, Tsunoda T, Deloukas P, Bird CP, Delgado M, Dermitzakis ET, Gwilliam R, Hunt S, Morrison J, Powell D, Stranger BE, Whittaker P, Bentley DR, Daly MJ, de Bakker PI, Barrett J, Chretien YR, Maller J, McCarroll S, Patterson N, Pe'er I, Price A, Purcell S, Richter DJ, Sabeti P, Saxena R, Schaffner SF, Sham PC, Varilly P, Altschuler D, Stein LD, Krishnan L, Smith AV, Tello-Ruiz MK, Thorisson GA, Chakravarti A, Chen PE, Cutler DJ, Kashuk CS, Lin S, Abecasis GR, Guan W, Li Y, Munro HM, Qin ZS, Thomas DJ, McVean G, Auton A, Bottolo L, Cardin N, Eyheramendy S, Freeman C, Marchini J, Myers S, Spencer C, Stephens M, Donnelly P, Cardon LR, Clarke G, Evans DM, Morris AP, Weir BS, Tsunoda T, Mullikin JC, Sherry ST, Feolo M, Skol A, Zhang H, Zeng C, Zhao H, Matsuda I, Fukushima Y, Macer DR, Suda E, Rotimi CN, Adebamowo CA, Ajayi I, Anigwu T, Marshall PA, Nkwojima C, Royal CD, Leppert MF, Dixon M, Peiffer A, Qiu R, Kent A, Kato K, Niikawa N, Adewole IF, Knoppers BM, Foster MW, Clayton EW, Watkin J, Gibbs RA, Belmont JW, Muzny D, Nazareth L, Sodergren E, Weinstock GM, Wheeler DA, Yakub I, Gabriel SB, Onofrio RC, Richter DJ, Ziaugra L, Birren BW, Daly MJ, Altschuler D, Wilson RK, Fulton LL, Rogers J, Burton J, Carter NP, Clee CM, Griffiths M, Jones MC, McLay K, Plumb RW, Ross MT, Sims SK, Willey DL, Chen Z, Han H, Kang L, Gdoubout M, Wallenburg JC, L'Archeveque P, Bellemare G, Saeki K, Wang H, An D, Fu H, Li Q, Wang Z, Wang R, Holden AL, Brooks LD, McEwen JE, Guyer MS, Wang VO, Peterson JL, Shi M, Spiegel J, Sung LM, Zacharia LF, Collins FS, Kennedy K, Jamieson R, Stewart J. A second generation human haplotype map of over 3.1 million SNPs. *Nature*. 2007;449:851–861.
  33. International HapMap C, Altschuler DM, Gibbs RA, Peltonen L, Altschuler DM, Gibbs RA, Peltonen L, Dermitzakis E, Schaffner SF, Yu F, Peltonen L, Dermitzakis E, Bonnen PE, Altschuler DM, Gibbs RA, de Bakker PI, Deloukas P, Gabriel SB, Gwilliam R, Hunt S, Inouye M, Jia X, Palotie A, Parkin M, Whittaker P, Yu F, Chang K, Hawes A, Lewis LR, Ren Y, Wheeler D, Gibbs RA, Muzny DM, Barnes C, Darvishi K, Hurler M, Korn JM, Kristiansson K, Lee C, McCarroll SA, Nemesh J, Dermitzakis E, Keinan A, Montgomery SB, Pollack S, Price AL, Soranzo N, Bonnen PE, Gibbs RA, Gonzaga-Jauregui C, Keinan A, Price AL, Yu F, Anttila V, Brodeur W, Daly MJ, Leslie S, McVean G, Moutsianas L, Nguyen H, Schaffner SF, Zhang Q, Ghorji MJ, McGinnis R, McLaren W, Pollack S, Price AL, Schaffner SF, Takeuchi F, Grossman SR, Shlyakhter I, Hostetter EB, Sabeti PC, Adebamowo CA, Foster MW, Gordon DR, Licinio J, Manca MC, Marshall PA, Matsuda I, Ngare D, Wang VO, Reddy D, Rotimi CN, Royal CD, Sharp RR, Zeng C, Brooks LD, McEwen JE. Integrating common and rare genetic variation in diverse human populations. *Nature*. 2010;467:52–58.
  34. Pistis G, Porcu E, Vrieze SI, Sidore C, Steri M, Danjou F, Busonero F, Mulas A, Zoledziwska M, Maschio A, Brennan C, Lai S, Miller MB, Marcelli M, Urru MF, Pitzalis M, Lyons RH, Kang HM, Jones CM, Angius A, Iacono WG, Schlessinger D, McGue M, Cucca F, Abecasis GR, Sanna S. Rare variant genotype imputation with thousands of study-specific whole-genome sequences: implications for cost-effective study designs. *Eur J Hum Genet*. 2014; doi: 10.1038/ejhg.2014.216 [Epub ahead of print].
  35. Janssens AC, Aulchenko YS, Elefante S, Borsboom GJJ, Steyberg EW, van Duijn CM. Predictive testing for complex diseases using multiple genes: fact or fiction? *Genet Med*. 2006;8:395–400.
  36. Janssens AC, van Duijn CM. Genome-based prediction of common diseases: advances and prospects. *Hum Mol Genet*. 2008;17:R166–R173.
  37. Talmud PJ, Hingorani AD, Cooper JA, Marmot MG, Brunner EJ, Kumari M, Kivimaki M, Humphries SE. Utility of genetic and non-genetic risk factors in prediction of type 2 diabetes: Whitehall II prospective cohort study. *BMJ*. 2010;340:b4838.
  38. Lysenko V, Laakso M. Genetic screening for the risk of type 2 diabetes: worthless or valuable? *Diabetes Care*. 2013;36(suppl 2):S120–S126.
  39. Paynter NP, Chasman DI, Pare G, Buring JE, Cook NR, Miletich JP, Ridker PM. Association between a literature-based genetic risk score and cardiovascular events in women. *JAMA*. 2010;303:631–637.
  40. Global Lipids Genetics C, Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, Ganna A, Chen J, Buchkovich ML, Mora S, Beckmann

- JS, Bragg-Gresham JL, Chang HY, Demirkan A, Den Hertog HM, Do R, Donnelly LA, Ehret GB, Esko T, Feitosa MF, Ferreira T, Fischer K, Fontanillas P, Fraser RM, Freitag DF, Gurdasani D, Heikkilä K, Hyppönen E, Isaacs A, Jackson AU, Johansson A, Johnson T, Kaakinen M, Kettunen J, Kleber ME, Li X, Luan J, Lyttikainen LP, Magnusson PK, Mangino M, Mihailov E, Montasser ME, Muller-Nurasyid M, Nolte IM, O'Connell JR, Palmer CD, Perola M, Petersen AK, Sanna S, Saxena R, Service SK, Shah S, Shungin D, Sidore C, Song C, Strawbridge RJ, Surakka I, Tanaka T, Teslovich TM, Thorleifsson G, Van den Herik EG, Voight BF, Volcik KA, Waite LL, Wong A, Wu Y, Zhang W, Absher D, Asiki G, Barroso I, Been LF, Bolton JL, Bonnycastle LL, Brambilla P, Burnett MS, Cesana G, Dimitriou M, Doney AS, Doring A, Elliott P, Epstein SE, Eyjolfsson GI, Gigante G, Goodarzi MO, Grallert H, Gravitto ML, Groves CJ, Hallmans G, Hartikainen AL, Hayward C, Hernandez D, Hicks AA, Holm H, Hung YJ, Illig T, Jones MR, Kaleebu P, Kastelein JJ, Khaw KT, Kim E, Klopp N, Komulainen P, Komari M, Langenberg C, Lehtimäki T, Lin SY, Lindstrom J, Loos RJ, Mach F, McArdle WL, Meisinger C, Mitchell BD, Muller G, Nagaraja R, Narisu N, Nieminen TV, Nsubuga RN, Olafsson I, Ong KK, Palotie A, Papamarkou T, Pomicino C, Pouta A, Rader DJ, Reilly MP, Ridker PM, Rivadeneira F, Rudan I, Ruokonen A, Samani N, Schanagl H, Seeley J, Silander K, Stancáková A, Stirrups K, Swift AJ, Tietel L, Uitterlinden AG, van Pelt LJ, Vedantam S, Wainwright N, Wijmenga C, Wild SH, Willemssen G, Wilsaard T, Wilson JF, Young EH, Zhao JH, Adair LS, Arveiler D, Assimes TL, Bandinelli S, Bennett F, Bochud M, Boehm BO, Boomsma DI, Borecki IB, Bornstein SR, Bovet P, Burnier M, Campbell H, Chakravarti A, Chambers JC, Chen YD, Collins FS, Cooper RS, Danesh J, Dedoussis G, de Faire U, Feranil AB, Ferrerres J, Ferrucci L, Freimer NB, Gieger C, Groop LC, Gudnason V, Gyllenstein U, Hamsten A, Harris TB, Hingorani A, Hirschhorn JN, Hofman A, Hovingh GK, Hsiung CA, Humphries SE, Hunt SC, Hveem K, Iribarren C, Jarvelin MR, Jula A, Kahonen M, Kaprio J, Kesäniemi A, Kivimäki M, Kooner JS, Koudstaal PJ, Krauss RM, Kuh D, Kuusisto J, Kyvik KO, Laakso M, Lakka TA, Lind L, Lindgren CM, Martin NG, Marz W, McCarthy MI, McKenzie CA, Meneton P, Metspalu A, Moilanen L, Morris AD, Munroe PB, Njolstad I, Pedersen NL, Power C, Pramstaller PP, Price JF, Psaty BM, Quertermous T, Rauramaa R, Saleheen D, Salomaa V, Sanghera DK, Saramies J, Schwarz PE, Sheu WH, Shuldiner AR, Siegbahn A, Spector TD, Stefansson K, Strachan DP, Tayo BO, Tremoli E, Tuomilehto J, Uusitupa M, van Duijn CM, Vollenweider P, Wallentin L, Wareham NJ, Whitfield JB, Wolffenbuttel BH, Ordovas JM, Boerwinkle E, Palmer CN, Thorsteinsdottir U, Chasman DI, Rotter JI, Franks PW, Ripatti S, Cupples LA, Sandhu MS, Rich SS, Boehnke M, Deloukas P, Kathiresan S, Mohlke KL, Ingelsson E, Abecasis GR. Discovery and refinement of loci associated with lipid levels. *Nat Genet.* 2013;45:1274–1283.
41. Hoggart CJ, Venturini G, Mangino M, Gomez F, Ascari G, Zhao JH, Teumer A, Winkler TW, Tsernikova N, Luan J, Mihailov E, Ehret GB, Zhang W, Lamparter D, Esko T, Mace A, Rueger S, Bochud PY, Barcella M, Dauvilliers Y, Benyamin B, Evans DM, Hayward C, Lopez MF, Franke L, Russo A, Heid IM, Salvi E, Vendantam S, Arking DE, Boerwinkle E, Chambers JC, Fiorito G, Grallert H, Guarrera S, Homuth G, Huffman JE, Porteous D; Generation Scotland C, LifeLines Cohort s, Consortium G, Moradpour D, Iranzo A, Hebebrand J, Kemp JP, Lammers GJ, Aubert V, Heim MH, Martin NG, Montgomery GW, Peraita-Adrados R, Santamaria J, Negro F, Schmidt CO, Scott RA, Spector TD, Strauch K, Volzke H, Wareham NJ, Yuan W, Bell JT, Chakravarti A, Kooner JS, Peters A, Matullo G, Wallaschowski H, Whitfield JB, Paccaud F, Vollenweider P, Bergmann S, Beckmann JS, Taft M, Hastie ND, Cusi D, Bochud M, Frayling TM, Metspalu A, Jarvelin MR, Scherag A, Smith GD, Borecki IB, Rousson V, Hirschhorn JN, Rivolta C, Loos RJ, Kutalik Z. Novel approach identifies SNPs in SLC2A10 and KCNK9 with evidence for parent-of-origin effect on body mass index. *PLoS Genet.* 2014;10:e1004508.
  42. Napoli C, D'Armiento FP, Mancini FP, Postiglione A, Witztum JL, Palumbo G, Palinski W. Fatty streak formation occurs in human fetal aortas and is greatly enhanced by maternal hypercholesterolemia. Intimal accumulation of low density lipoprotein and its oxidation precede monocyte recruitment into early atherosclerotic lesions. *J Clin Invest.* 1997;100:2680–2690.
  43. Neary RH, Kilby MD, Kumpatula P, Game FL, Bhatnagar D, Durrington PN, O'Brien PM. Fetal and maternal lipoprotein metabolism in human pregnancy. *Clin Sci.* 1995;88:311–318.
  44. Napoli C, Glass CK, Witztum JL, Deutsch R, D'Armiento FP, Palinski W. Influence of maternal hypercholesterolemia during pregnancy on progression of early atherosclerotic lesions in childhood: fate of early lesions in children (FELIC) study. *Lancet.* 1999;354:1234–1241.
  45. Kilby MD, Neary RH, Mackness MI, Durrington PN. Fetal and maternal lipoprotein metabolism in human pregnancy complicated by type I diabetes mellitus. *J Clin Endocrinol Metab.* 1998;83:1736–1741.
  46. Dabelea D, Hanson RL, Lindsay RS, Pettitt DJ, Imperatore G, Gabir MM, Roumain J, Bennett PH, Knowler WC. Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. *Diabetes.* 2000;49:2208–2211.
  47. Dabelea D. The predisposition to obesity and diabetes in offspring of diabetic mothers. *Diabetes Care.* 2007;30(suppl 2):S169–S174.
  48. Oken E, Levitan EB, Gillman MW. Maternal smoking during pregnancy and child overweight: systematic review and meta-analysis. *Int J Obes.* 2008;32:201–210.
  49. Wright CS, Rifas-Shiman SL, Rich-Edwards JW, Taveras EM, Gillman MW, Oken E. Intrauterine exposure to gestational diabetes, child adiposity, and blood pressure. *Am J Hypertens.* 2009;22:215–220.
  50. Dominguez-Salas P, Moore SE, Baker MS, Bergen AW, Cox SE, Dyer RA, Fulford AJ, Guan Y, Laritsky E, Silver MJ, Swan GE, Zeisel SH, Innis SM, Waterland RA, Prentice AM, Hennig BJ. Maternal nutrition at conception modulates DNA methylation of human metastable epialleles. *Nat Commun.* 2014;5:3746.
  51. Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet.* 1986;1:1077–1081.
  52. Paneth N, Susser M. Early origin of coronary heart disease (the “Barker hypothesis”). *BMJ.* 1995;310:411–412.
  53. Jorgensen AB, Frikke-Schmidt R, Nordestgaard BG, Tybjaerg-Hansen A. Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *N Engl J Med.* 2014;371:32–41.
  54. Tg, Hdl Working Group of the Exome Sequencing Project NHL, Blood I, Crosby J, Peloso GM, Auer PL, Crosslin DR, Stitzel NO, Lange LA, Lu Y, Tang ZZ, Zhang H, Hindy G, Masca N, Stirrups K, Kanoni S, Do R, Jun G, Hu Y, Kang HM, Xue C, Goel A, Farrall M, Duga S, Merlini PA, Asselta R, Girelli D, Olivieri O, Martinelli N, Yin W, Reilly D, Speliotes E, Fox CS, Hveem K, Holmen OL, Nikpay M, Farlow DN, Assimes TL, Franceschini N, Robinson J, North KE, Martin LW, DePristo M, Gupta N, Escher SA, Jansson JH, Van Zuydam N, Palmer CN, Wareham N, Koch W, Meitinger T, Peters A, Lieb W, Erbel R, König IR, Kruppa J, Degenhardt F, Gottesman O, Bottinger EP, O'Donnell CJ, Psaty BM, Ballantyne CM, Abecasis G, Ordovas JM, Melander O, Watkins H, Orho-Melander M, Ardisino D, Loos RJ, McPherson R, Willer CJ, Erdmann J, Hall AS, Samani NJ, Deloukas P, Schunkert H, Wilson JG, Kooperberg C, Rich SS, Tracy RP, Lin DY, Altshuler D, Gabriel S, Nickerson DA, Jarvik GP, Cupples LA, Reiner AP, Boerwinkle E, Kathiresan S. Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N Engl J Med.* 2014;371:22–31.